SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If the information provided is not</u> complete, correct, or legible, the authorization process can be delayed.

Ophthalmic Immunomodulator Drugs

Drug Requested: (Check below the drug that applies)

PREFERRED (must be tried and failed FIRST)		
	Restasis [®] (cyclosporine)	□ Xiidra [®]
Non-Preferred		
	Cequa [™] (cyclosporine)	□ Eysuvis [®] (loteprednol)
	cyclosporine	□ Miebo [™]
	Restasis Multidose [®] (cyclosporine)	□ Tyrvaya [™] (varenicline tartrate) Nasal Spray
	Verkazia [®] (cyclosporine) *Trial and failure of preferred agents do not apply	

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Date of Birth:
Date:
Fax Number:
ed if incomplete.
Length of Therapy:
CD Code, if applicable:
Date weight obtained:

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

□ Cequa[™] (cyclosporine), cyclosporine, Miebo[™], Restasis Multidose[®] (cyclosporine), Tyrvaya[™] (varenicline tartrate) Nasal Spray

□ Member has a trial and failure of <u>TWO (2)</u> preferred alternatives: (Check all that apply)

 \Box Restasis[®] (cyclosporine)

□ Xiidra[®]

Eysuvis[®] (loteprednol) Quantity Limit: 1 bottle per 3 months

□ Member has a diagnosis of dry eye disease

AND

□ Prescriber attest to utilizing Eysuvis[®] for short-term treatment (up to 2 weeks of therapy)

AND

□ Member has a trial and failure of <u>TWO (2)</u> preferred alternatives:

□ Restasis [®] (cyclosporine)	□ Xiidra [®]
--	-----------------------

Verkazia[®] (cyclosporine) (Routine PDL criteria does not apply) Quantity Limit: 120 single-dose vials per 30 days 1 bottle per 3 months

- □ Member is at least 4 years of age
- **D** The provider is an ophthalmologist or an optometrist in consultation with an ophthalmologist
- □ Member has a diagnosis of severe vernal keratoconjunctivitis
- □ Member has trial and failure, contraindication or intolerance to at least **one** of the following:
 - **D** Topical ophthalmic "dual action" mast cell stabilizer and antihistamine (e.g., olopatadine, azelastine)
 - □ Topical ophthalmic mast cell stabilizers (e.g., cromolyn)

(Continued on next page)

MEDICAL NECESSITY: Provide clinical evidence that the <u>**PREFERRED</u>** drugs will not provide adequate benefit.</u>

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. *<u>Previous therapies will be verified through pharmacy paid claims or submitted chart notes.</u>*