

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Ophthalmic Immunomodulator Drugs

Drug Requested: (Check below the drug that applies)

PREFERRED (must be tried and failed FIRST)	
<input type="checkbox"/> Restasis®/Restasis Multidose® (cyclosporine)	<input type="checkbox"/> Xiidra® (cyclosporine)
Non-Preferred	
<input type="checkbox"/> Cequa™ (cyclosporine)	<input type="checkbox"/> Eysuvis® (loteprednol)
<input type="checkbox"/> cyclosporine	<input type="checkbox"/> Miebo™
<input type="checkbox"/> Tyrvaya™ (varenicline tartrate) Nasal Spray	<input type="checkbox"/> Verkazia® (cyclosporine) *Trial and failure of preferred agents do not apply

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Cequa™ (cyclosporine), cyclosporine, Miebo™, Tyrvaya™ (varenicline tartrate) Nasal Spray

- Member has a trial and failure of **TWO (2)** preferred alternatives: **(Check all that apply)**

Restasis®/Restasis Multidose® (cyclosporine)

Xiidra® (cyclosporine)

Eysuvis® (loteprednol)

Quantity Limit: 1 bottle per 3 months

- Member has a diagnosis of dry eye disease

AND

- Prescriber attest to utilizing Eysuvis® for short-term treatment (up to 2 weeks of therapy)

AND

- Member has a trial and failure of **TWO (2)** preferred alternatives:

Restasis®/Restasis Multidose® (cyclosporine)

Xiidra® (cyclosporine)

Verkazia® (cyclosporine) (Routine PDL criteria does not apply)

Quantity Limit: 120 single-dose vials per 30 days

- Member is at least 4 years of age

AND

- The provider is an ophthalmologist or an optometrist in consultation with an ophthalmologist

AND

- Member has a diagnosis of severe vernal keratoconjunctivitis

AND

- Member has trial and failure, contraindication or intolerance to one of the following:

- Topical ophthalmic “dual action” mast cell stabilizer and antihistamine (e.g., olopatadine, azelastine)

OR

- Topical ophthalmic mast cell stabilizers (e.g., cromolyn)

MEDICAL NECESSITY: Provide clinical evidence that the **PREFERRED** drugs will not provide adequate benefit.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****