SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed.</u>

Drug Requested: Primaxin® IV (cilastatin sodium/imipenem) J0743 (Medical)

MEMBER & PRESCRIBER INFORMATIO	N: Authorization may be delayed if incomplete.	
Member Name:		
Member Optima #:		
Prescriber Name:		
Prescriber Signature:	Date:	
Office Contact Name:		
Phone Number:	Fax Number:	
NPI #:		
DRUG INFORMATION: Authorization may be d	lelayed if incomplete.	
Drug Form/Strength:		
Dosing Schedule:	Length of Therapy:	
Diagnosis:	ICD Code, if applicable:	
Weight (if applicable):	Date weight obtained:	
☐ Standard Review. In checking this box, the timeframe or the member's ability to regain maximum function a	v 1	
CLINICAL CRITERIA: Check below all that appropriate support each line checked, all documentation, including provided or request may be denied.		
Length of Authorization: Date of Service (14	days)	
□ New Start		

(Continued on next page)

	Member has ONE of the following diagnoses:		
		Lower respiratory tract infections	
		Urinary tract infections	
		Intra-abdominal infections	
		Gynecologic infections	
		Bone and joint infections	
		Skin and skin structure infections	
		Endocarditis	
	Provider has submitted lab cultures from current hospital admission or office visit collected within the last 7 days		
	□ Provider must submit chart notes documenting trial and failure of at least <u>TWO</u> of the following or IV preferred antibiotics within the last 14 days specific to the applicable indication for use:		
		Lower respiratory tract infections – ceftriaxone, azithromycin, cefepime, doxycycline, and levofloxacin	
		Urinary tract infections – nitrofurantoin, cefdinir, cephalexin, amoxicillin, amoxicillin-clavulanate, ciprofloxacin, levofloxacin, trimethoprim-sulfamethoxazole, and fosfomycin	
		Intra-abdominal infections – ciprofloxacin, levofloxacin, ceftriaxone, cefazolin, cefepime, piperacillin-tazobactam, trimethoprim-sulfamethoxazole, ertapenem, imipenem-cilastatin, and meropenem	
		Gynecologic infections – nitrofurantoin, cefdinir, cephalexin, amoxicillin, amoxicillin-clavulanate, ciprofloxacin, levofloxacin, trimethoprim-sulfamethoxazole, and fosfomycin	
		Bone and joint infections – vancomycin, nafcillin, oxacillin, cefazolin, ceftriaxone, daptomycin, cipro, levofloxacin, ceftazidime, and ertapenem	
		Skin and skin structure infections – cephalexin, dicloxacillin, cefazolin, ceftriaxone, piperacillin-tazobactam, vancomycin, trimethoprim-sulfamethoxazole, doxycycline, clindamycin, and ciprofloxacin	
		Endocarditis – vancomycin, ceftriaxone, gentamicin, daptomycin, cefepime, piperacillin-tazobactam tobramycin, meropenem	
ength of Authorization: Date of Service			
C	on	tinuation of therapy following inpatient administration	
	Me	ember has ONE of the following diagnoses:	
		Lower respiratory tract infections	
		Urinary tract infections	
		Intra-abdominal infections	
		Gynecologic infections	
		Bone and joint infections	
		Skin and skin structure infections	
		Endocarditis	

(Continued on next page)

PA Primaxin IV (Medical)(Medicaid) (Continued from previous page)

 □ Member is currently on Primaxin for more than 72 hours inpatient (progress notes must be submitted) □ Provider has submitted lab culture sensitivity results retrieved during inpatient admission which shows resistance to <u>ALL</u> preferred antibiotics except for Primaxin (sensitive)
Medication being provided by: Please check applicable box below.
□ Location/site of drug administration:
NPI or DEA # of administering location:
<u>OR</u>
□ Specialty Pharmacy
For urgent reviews: Practitioner should call Optima Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Optima's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.
Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. *Previous therapies will be verified through pharmacy paid claims or submitted chart notes.*