SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Sickle Cell Disease Drugs

Drug Requested: (Please select drug below)

PREFERRED MEDICATIONS (Does not require Prior Authorization for FDA approved ages)		
□ Droxia®	□ Endari™	
	-PREFERRED MEDICATIONS Require prior authorization)	
□ Adakveo IV®	□ Siklos®	
MEMBER & PRESCRIBER I	NFORMATION: Authorization may be delayed if incomplete.	
Member Name:		
Member Sentara #:	Date of Birth:	
Prescriber Name:		
Prescriber Signature:	Date:	
Office Contact Name:		
Phone Number:		
NPI #:		
DRUG INFORMATION: Auth	orization may be delayed if incomplete.	
Drug Name/Form/Strength:		
Dosing Schedule:	Length of Therapy:	
Diagnosis:		
Weight (if applicable):	Date weight obtained:	

(Continued on next page)

	support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.					
Init	ial Approval: 6 months					
1.	Is the drug being prescribed by or in consultation with an oncologist, hematologist or sickle cell specialist? ☐ Yes ☐ No					
2.	Does the patient have a diagnosis of Sickle Cell Disease presenting as one of following (HbSS, HbSC, HbS β ⁰ -thalassemia, or HbS β ⁺ -thalassemia)?					
	□ Yes □ No					
3.	3. Is the medication dose proper for the patient's age or other conditions affecting the dose, according to t product package insert approved by the FDA?					
	□ Yes □ No					
For A	Adakveo®					
4.	Has the member had an insufficient response to a minimum 3-month trial of hydroxyurea (unless contraindicated or intolerant)?					
	□ Yes □ No					
5.	Patient has experienced TWO or more vaso-occlusive crises (VOC) in the previous year despite adherence to hydroxyurea therapy?					
	□ Yes □ No					

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To

For Siklos®

6. Is the member between 2 to 17 years of age?

□ Yes □ No

<u>Reauthorization Approval</u>: 1 year. All criteria must be checked for approval. To support each line checked, all documentation (lab results, diagnostics, and/or chart notes) must be provided or request may be denied.

1.	Does the r	 tinue to meet the above criteria?
2.	Does the r	e disease response improvement with treatment?

For Adakveo®

1. Is the member's response compared to pre-treatment baseline evidenced by a decrease in the frequency of vaso-occlusive crises (VOC) necessitating treatment, reduction in number or duration of hospitalizations, and/or reduction in severity of VOC?

□ Yes □ No

List pharmaceutical drugs attempted and outcome:		
Medical necessity: Provide clinical evidence that the <u>PREFERRED</u> drug(s) will not provide adequate benefit.		
Medication being provided by a Specialty Pharmacy - PropriumRx		

^{**}Use of samples to initiate therapy does not meet step edit/preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *