SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

<u>Drug Requested</u>: Gamifant® (emapalumab-lzsg) – HLH/MAS (J9210) MEDICAL

The Sentara Health Plans Oncology Program is administered by OncoHealth

- ❖ For any oncology indications, the most efficient way to submit a prior authorization request is through the OncoHealth OneUM Provider Portal at https://oneum.oncohealth.us. Fax to 1-800-264-6128. OncoHealth can also be contacted by Phone: 1-888-916-2616.
- ❖ Commercial customers <u>NOT</u> enrolled in the OncoHealth program, please fax requests to Sentara Health plans at fax number 1-800-750-9692.

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete. Member Name: _____ Member Sentara #: Date of Birth: Prescriber Name: Prescriber Signature: Date: Office Contact Name: Phone Number: Fax Number: NPI #: **DRUG INFORMATION:** Authorization may be delayed if incomplete. Drug Name/Form/Strength: Dosing Schedule: Length of Therapy: Diagnosis: ______ ICD Code, if applicable: _____ Date weight obtained: _____ Weight (if applicable): □ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

(Continued on next page)

Max Units (per dose and over time) [HCPCS Unit]: 9250 billable units per 30 days; 1 mg = 1 billable unit

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.				
	Diagnosis: Hemophagocytic Lymphohistiocytosis (HLH)/Macrophage Activation Syndrome (MAS) in known or suspected Still's disease			
Init	ial Authorization: 30 days			
	Member has a definitive diagnosis of HLH/MAS as indicated by BOTH of the following (submit documentation):			
	□ Ferritin >684 ng/mL			
	☐ At least 2 of the following (check all that apply):			
	□ Platelet count $\leq 181 \times 10^9 / L$			
	\Box AST >48 U/L			
	☐ Triglycerides >156 mg/dL			
	☐ Fibrinogen levels ≤360 mg/dL			
	Member has known or suspected diagnosis of Still's disease, including systemic Juvenile Idiopathic Arthritis (sJIA) or Adult Onset Still's Disease (AOSD)			
	Member must meet ONE of the following (submit documentation):			
	☐ Member has had an inadequate response or intolerance to high-dose intravenous (IV) glucocorticoids (currently or last 30 days)			
	☐ Member has recurrent MAS			
	Member has been evaluated and screened for the presence of latent tuberculosis (TB) infection prior to initiating treatment and will receive ongoing monitoring for the presence of TB during treatment			
	Providers will monitor and consider prophylaxis in patients for Herpes Zoster, <i>Pneumocystis Jirovecii</i> , and fungal infections			
	Member does <u>NOT</u> have an active infection, including clinically important localized infections that are favored by interferon-gamma neutralization (e.g., infections caused by mycobacteria, Histoplasma Capsulatum)			
	Medication must NOT be administered concurrently with live or live attenuated vaccines			
supp	uthorization: 30 days. Check below all that apply. All criteria must be met for approval. To ort each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be ided or request may be denied.			
	Member continues to require therapy for treatment of HLH/MAS			

(Continued on next page)

(0 . 10	•	•	
(Continued f	rom	previous	page

- ☐ Member must meet <u>ONE</u> of the following:
 - ☐ Member experienced a complete response (CR) as evidenced by the following:
 - □ Clinical resolution of MAS signs and symptoms (a visual analogue scale (VAS), of ≤1 cm [range 0 to 10 cm])
 - ☐ Member meets ALL the following laboratory parameter endpoints:
 - □ WBC count and platelet count above the lower limit of normal (LLN)
 - □ LDH, AST and ALT below 1.5 times the upper limit of normal (ULN)
 - ☐ Fibrinogen >100 mg/dL
 - ☐ Ferritin levels decreased ≥80% from values at screening or baseline (whichever initial value was higher) or < 2000 ng/mL, whichever was lower
- ☐ Member has had unsatisfactory improvement in clinical condition, as assessed by a healthcare provider and requires dose escalation (up to the maximum dose and frequency specified in the Dosage/Administration table below)

Treatment Day	Gamifant Dose	Dose Adjustments	
Day 1	Initial Dose of 6 mg/kg	If unsatisfactory improvement in clinical condition, as assessed by a healthcare provider, the dose of Gamifant may be increased to:	
Days 4-16	3 mg/kg every 3 days for 5 doses	A maximum cumulative dose of 10 mg/kg over 3 days	
Day 19 onwards	3 mg/kg twice per week (i.e., every 3 to 4 days)	AND the frequency may be increased to: • Every 2 days or once daily After the patient's clinical condition has improved, consider decreasing the dose to the previous level and assess whether clinical response is maintained. If the clinical condition is not stabilized while receiving the maximum dosage, consider discontinuing Gamifant.	

- ☐ Member has experienced an absence of unacceptable toxicity from the drug (e.g., serious infections (including mycobacteria, Herpes Zoster virus, and Histoplasma Capsulatum)
- ☐ Member is receiving ongoing monitoring for adenovirus, EBV, and CMV viruses as clinically indicated

(Continued on next page)

Medication being provided by: Please check applicable box below.					
	Location/site of drug administration:				
	NPI or DEA # of administering location:				
	<u>OR</u>				
	Specialty Pharmacy				
a star of ur	argent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe indard review would subject the member to adverse health consequences. Sentara Health Plan's definition gent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's y to regain maximum function.				
	Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. revious therapies will be verified through pharmacy paid claims or submitted chart notes.*				