

OPTIMA HEALTH PLAN

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization may be delayed.**

Gastrointestinal (GI) Motility Drugs

Drug Requested: (select one drug below)		
Non-Preferred		
<input type="checkbox"/> Ibsrela[®] (tenapanor)	<input type="checkbox"/> Motegrity[®] (prucalopride)	<input type="checkbox"/> Relistor[®] (methylnaltrexone bromide)
<input type="checkbox"/> Trulance[®] (plecanatide)	<input type="checkbox"/> Zelnorm[™] (tegaserod)	

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Optima #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

(Continued on next page)

☐ **Approval of Motegrity® for diagnosis of Chronic Idiopathic Constipation (CIC)**

- ☐ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to Linzess®

☐ **Approval of Zelnorm™**

- ☐ Diagnosis of **Irritable Bowel Syndrome with Constipation (IBS-C)**

AND

- ☐ Member is < 65 years of age with no history of ischemic cardiovascular disease and has no more than one CVD risk factor. CVD risk factors are defined as active smoking, current hypertension/history of antihypertensive treatment, current hyperlipidemia/history of lipid lowering medication, history of diabetes mellitus, age >55 years, or obesity (BMI >30 kg/m²)

AND

- ☐ Provider attests that member does **NOT** have any of the following contraindications to therapy:
- History of myocardial infarction (MI), stroke, transient ischemic attack (TIA), or angina
 - History of ischemic colitis or other forms of intestinal ischemia
 - Severe renal impairment (eGFR < 15 mL/min/1.73 m²) or end-stage renal disease
 - Moderate and severe hepatic impairment (Child-Pugh B or C)
 - History of bowel obstruction, symptomatic gallbladder disease, suspected sphincter of Oddi dysfunction, or abdominal adhesions

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

AND

(Continued on next page)

- ☐ Member has had trial and failure, contraindication, or intolerance to Linzess®

☐ Approval of Trulance® for diagnosis of Irritable Bowel Syndrome with Constipation (IBS-C) or Chronic Idiopathic Constipation (CIC)

- ☐ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to Linzess®

☐ Approval of Ibsrela® for diagnosis of Irritable Bowel Syndrome with Constipation (IBS-C)

- ☐ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to Linzess®

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to Trulance® (**requires prior authorization**)

☐ Approval of Relistor®

Recommended Dosing:

Weight of Adult Patient	Subcutaneous Dose	Injection Volume
Less than 38kg	0.15 mg/kg	See below
38kg to less than 62 kg	8mg	0.4 mL
62kg to 114kg	12mg	0.6 mL
More than 114kg	0.15 mg/kg	See below

Calculate injection volume by multiplying member weight in kilograms by 0.0075, then round up the volume to the nearest 0.1 mL

(Continued on next page)

- ☐ Select **ONE** of the following:

- ☐ Member has a diagnosis of opioid-induced constipation (OIC) with advanced illness or pain caused by active cancer who require opioid dosage escalation for palliative care
- ☐ Member has a diagnosis of opioid-induced constipation (OIC) with chronic non-cancer pain, including members with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation

AND

- ☐ Member has been on an opioid within the last 60 days of prior authorization request but not less than 4 weeks. Provider please note: Members receiving opioids for less than 4 weeks may be less responsive to Relistor[®]

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX [®])
------------------------------------	--

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza[®])

AND

- ☐ Member has had trial and failure, contraindication, or intolerance to both Movantik[®]
AND Symproic[®]

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****