

SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-305-2331**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

Drug Requested: Orencia® (abatacept) (J-0129) (IV INFUSION ONLY) (Medical)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

| DIAGNOSIS | Recommended Quantity |
|---|--|
| Moderately to severely active Rheumatoid Arthritis (RA) | INTRAVENOUS <ul style="list-style-type: none"> Weight <60kg: Four vials in initial 28 days; Two vials per 28 days Weight 60-100kg: Six vials in initial 28 days; Three vials per 28 days Weight >100kg: Eight vials in initial 28 days; Four vials per 28 days |

(Continued on next page)

| DIAGNOSIS | Recommended Quantity |
|--|---|
| Psoriatic arthritis (PsA) | INTRAVENOUS <ul style="list-style-type: none"> • Weight <60kg: Four 250mg vials initial 28 days; Two vials per 28 days after induction • Weight 60-100kg: Six vials in initial 28 days; Three vials per 28 days after induction • Weight >100kg: Eight vials initial 28 days; Four vials per 28 days after induction |
| Juvenile Idiopathic Arthritis (JIA) in members 2 years and older | INTRAVENOUS <ul style="list-style-type: none"> • ≥ 6 y/o (subcutaneous formulation indicated for 2 years and older) • Weight <75kg: 10mg/kg every 28 days (3 vial) • Weight >75kg: Follow adult Rheumatoid Arthritis dosing above (not to exceed max dose 1000mg) |
| Prophylaxis of acute graft versus host disease (aGVHD) | INTRAVENOUS <ul style="list-style-type: none"> • Patients 2 to less than 6 years old, administer a 15 mg/kg dose as a 60-minute infusion on the day before transplantation, followed by a 12 mg/kg dose as a 60-minute infusion on Day 5, 14, and 28 after transplant • Patients 6 years and older, 10 mg/kg dose (maximum dose 1,000 mg) as a 60-minute infusion day before transplantation, followed by a dose on Day 5, 14, and 28 after transplant |

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Moderate to severe Active Rheumatoid Arthritis (RA)

- Member is 18 years of age or older
- Member has been diagnosed with **moderate to severe rheumatoid arthritis (RA)**
- Trial and failure of, contraindication, or adverse reaction to methotrexate
- Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**)

| | |
|---|--|
| <input type="checkbox"/> auranofin | <input type="checkbox"/> sulfasalazine |
| <input type="checkbox"/> azathioprine | <input type="checkbox"/> leflunomide |
| <input type="checkbox"/> hydroxychloroquine | <input type="checkbox"/> other: _____ |

(Continued on next page)

PA Orencia IV (MEDICAL) (Medicaid)

(Continued from previous page)

- Member has tried and failed **BOTH** of the preferred drugs below:

| | |
|---|--|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd) | <input type="checkbox"/> Enbrel [®] |
|---|--|

Polyarticular Juvenile Idiopathic Arthritis (pJIA)

- Member is 2 years of age or older
- Member has been diagnosed with **moderate to severe active polyarticular Juvenile Idiopathic Arthritis (pJIA)**
- Trial and failure of **BOTH** of the preferred below:

| | |
|---|--|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd) | <input type="checkbox"/> Enbrel [®] |
|---|--|

Active Psoriatic Arthritis (PsA)

- Member is 2 years of age or older
- Member has been diagnosed with **moderate to severe active Psoriatic Arthritis (PsA)**
- Member has tried and failed **TWO (2)** of the preferred drugs below:

| | | |
|--|--|--|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd) | <input type="checkbox"/> Enbrel [®] | <input type="checkbox"/> Pyzchiva [®] syringe/vial OR Starjemza [™] (Requires trial and failure of a preferred TNF-alpha inhibitor) |
|--|--|--|

Prophylaxis of acute graft versus host disease (aGVHD)

- Member has a diagnosis of prophylaxis of **acute graft versus host disease (aGVHD)**
- Member is 2 years of age and older undergoing hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor
- The medication is used in combination with a calcineurin inhibitor and methotrexate

(Continued on next page)

Medication being provided by: Please check applicable box below.

- Location/site of drug administration: _____
NPI or DEA # of administering location: _____

OR

- Physician's office **OR** Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****