SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Taltz® SQ (ixekizumab)

every 4 weeks

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.		
Member Name:		
Member Sentara #:		
Prescriber Name:		
Prescriber Signature:	Date:	
Office Contact Name:		
Phone Number:	Fax Number:	
DEA OR NPI #:		
DRUG INFORMATION: Authorization may be	delayed if incomplete.	
Drug Form/Strength:		
Dosing Schedule:		
Diagnosis:	ICD Code, if applicable:	
Weight:	Date:	
CLINICAL CRITERIA: Check below all that ap support each line checked, all documentation, including provided or request may be denied. Check the diagnosis	lab results, diagnostics, and/or chart notes, must be	
□ Diagnosis: Moderate-to-Severe Chronic P	Plaque Psoriasis	
Dosing:		
Adults: SubQ: Initial: 160 mg once, followed by 80 mg every 4weeks	0 mg at weeks 2, 4, 6, 8, 10, and 12. Maintenance:	
<u>Pediatrics</u> :		
Children ≥ 6 years and Adolescents <18 years:		
• < 25 kg: SubQ: 40 mg once, followed by 20	•	
 25 to 50 kg: SubQ: 80 mg once, followed by > 50 kg: SubQ: 160 mg once (administered at the subQ: 160 mg once) 	as 2 separate 80 mg injections), followed by 80 mg	

(Continued on next page)

	Member is ≥ 6 years of age and has a diagnosis of moderate-to-severe plaque psoriasis		
	Prescribed by or in consultation with a Dermatologist		
	Member tried and failed at least ONE of either Phototherapy or Alternative Systemic Therapy for at least three (3) months (check each tried below):		
	□ Phototherapy: □ UV Light Therapy □ NB UV-B □ PUVA	□ Alternative Systemic Therapy: □ Oral Medications □ acitretin □ methotrexate □ cyclosporine	
□ D	iagnosis: Active Psoriatic Arthritis		
Dosin	ig: SubQ: 160 mg once, followed by 80 mg every	ł weeks	
- -	Member has a diagnosis of active psoriatic arthri Prescribed by or in consultation with a Rheumato Member tried and failed at least ONE of the follow □ methotrexate oral or SQ 15-25 mg/week □ leflunomide oral 20 mg/day □ sulfasalazine oral 2-3 g/day		
□ D	iagnosis: Active Ankylosing Spondylitis		
Dosing: SubQ: 160 mg once, followed by 80 mg every 4 weeks			
<u> </u>	Member has a diagnosis of active ankylosing sport Prescribed by or in consultation with a Rheumato Member tried and failed, has a contraindication, or	logist	
□ Diagnosis: Active Non-radiographic Axial Spondyloarthritis			
Dosing: SubQ: 80 mg every 4 weeks			
<u> </u>	Member has a diagnosis of active non-radiograph Prescribed by or in consultation with a Rheumato	•	

(Continued on next page)

PA Taltz SQ (CORE) (Continued from previous page)

	Member has at least ONE of the following objective signs of inflammation:	
	☐ C-reactive protein [CRP] levels above the upper limit of normal	
	□ Sacroiliitis on magnetic resonance imaging [MRI] (indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints)	
	Member tried and failed, has a contraindication, or intolerance to TWO NSAIDs	
Medication being provided by a Specialty Pharmacy - PropriumRx		

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *