

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization will be delayed.**

Antibiotics-Inhaled

Drug Requested: (Check below all that apply)

PREFERRED	
<input type="checkbox"/> Bethkis [®] 300 mg/4mL (QL, AG)	<input type="checkbox"/> Kitabis [®] Pak 300 mg/5mL (QL, AG)
<input type="checkbox"/> Tobi Podhaler [®] (QL, AG, SE)	<input type="checkbox"/> tobramycin inhalation neb soln 300mg/5mL (generic Tobi [®] inhalation) (QL, AG)
Non-Preferred	
<input type="checkbox"/> Arikayce [®] (QL, PA required)	<input type="checkbox"/> Cayston [®] (QL, AG)
<input type="checkbox"/> Tobi [®] inhalation neb soln 300mg/5mL (QL, AG)	<input type="checkbox"/> tobramycin inhalation (generic Bethkis [®]) (QL, AG)
<input type="checkbox"/> tobramycin Pak (generic Kitabis [®] Pak) (QL, AG)	

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

(Continued on next page)

Quantity Limits		
Arikayce = 590 mg/8.4 mL (28 vials)/ 28 days	Bethkis® = 224 mL (56 amps)/ 28 days	Cayston® = 84 mL (56 amps)/ 28 days
Kitabis® Pak = 280 mL(56 amps)/ 28 days	Tobi Podhaler® = 224 capsule/ 28 days	Tobi® inhalation neb = 280mL (56 amps)/ 28 days
tobramycin = 280mL (56 amps)/ 28 days		

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Minimum age for use is 6 years for all tobramycin inhalation nebulizer solution (Bethkis®, Kitabis™ Pak, Tobi® and Tobi Podhaler®) and 7 years for Cayston®.

- **Tobi Podhaler®:** clinical reason why ONE (1) of the **PREFERRED** tobramycin inhalation nebulizer solutions cannot be used.

******Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.******
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****