# SENTARA COMMUNITY PLAN (MEDICAID)

## MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed.</u>

## The Sentara Health Plans Oncology Program is administered by OncoHealth

❖ For any oncology indications, the most efficient way to submit a prior authorization request is through the OncoHealth OneUM Provider Portal at <a href="https://oneum.oncohealth.us">https://oneum.oncohealth.us</a>. Fax to 1-800-264-6128. OncoHealth can also be contacted by Phone: 1-888-916-2616.

□ Riabni <sup>™</sup> (rituximab-arrx) (Q5123)	□ Rituxan® (rituximab) (J9312)		
□ Ruxience <sup>™</sup> (rituximab-pvvr) (Q5119)	□ Truxima® (rituximab-abbs) (Q5115)		
MEMBER & PRESCRIBER INFORMA	<b>TION:</b> Authorization may be delayed if incomplete.		
Member Name:			
Member Sentara #:	Date of Birth:		
Prescriber Name:			
Prescriber Signature:			
Office Contact Name:			
Phone Number:			
NPI #:			
DRUG INFORMATION: Authorization may			
Drug Name/Form/Strength:			
sing Schedule: Length of Therapy:			
gnosis: ICD Code, if applicable:			
Weight (if applicable):	Date weight obtained:		
☐ Standard Review. In checking this box, the time	frame does not jeopardize the life or health of the member		

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

or the member's ability to regain maximum function and would not subject the member to severe pain.

<b>DIAGNOSES:</b> Check below one of the diagnoses that applies to qualify.					
□ Rheumatoid Arthritis (RA)					
	Prescriber is a rheumatologi	st			
	<u>AND</u>				
	Member has a diagnosis of moderate- to-severe rheumatoid arthritis				
	AND				
	☐ Trial and failure of at least three (3) months of methotrexate therapy				
	AND				
	Trial and failure of TWO (2	c) of the preferred medical	biologics:		
	☐ Humira <sup>®</sup>	□ Enbrel <sup>®</sup>		1 Infliximab	
<b>-</b> (	Granulomatosis with Pol	yangiitis OR Microsco	opic Polyan	giitis	
INI	TIAL THERAPY [Ruxi	ence, Truxima, Rituxa	ın, Riabni]		
	☐ Prescriber is a rheumatologist or nephrologist				
	<u>AND</u>				
	☐ Member has a diagnosis of moderate- to-severe granulomatosis with polyangiitis				
	<u>AND</u>				
	Member will receive concur	rent therapy with corticost	eroids		
	AND				
	Member failed cyclophosph	amide therapy			
	<u>OR</u>				
	Member has a contraindicat	ion to cyclophosphamide the	herapy:		

(Continued on next page)

<b>-</b> (	Granulomatosis with Polyangiitis OR Microscopic Polyangiitis
MA	INTENANCE THERAPY [Ruxience, Truxima, Rituxan, Riabni]
	Prescriber is a rheumatologist or nephrologist
	AND
	Induction occurred at least 4 months prior
	AND
	Total duration of treatment does not exceed 24 months
	AND
	Member failed methotrexate or azathioprine therapy
	<u>OR</u>
	Member has a contraindication to methotrexate or azathioprine therapy:
	euromyelitis Optica Spectrum Disorder (NMOSD) [Ruxience, Truxima, Rituxan, liabni]
	Prescribing physician must be a neurologist
	AND
	Member must be 18 years of age or older
	AND
	Provider must submit medical records (e.g., chart notes) confirming the member has a diagnosis of Neuromyelitis Optica Spectrum Disorder (NMOSD) based on the following:
	☐ Member was found to be seropositive for aquaporin-4 (AQP4) IgG antibodies; <b>AND</b>
	☐ Member has at least one core clinical characteristic (see chart below)
	OR
	☐ Member was found to be seronegative for AQP-4 IgG antibodies OR has unknown AQP-4- IgG status
	AND
	☐ Member has at least two core clinical characteristics occurring as a result of one or more clinical attacks (see chart below)
	AND

#### **AND**

- ☐ Member experienced <u>ALL</u> the following:
  - ☐ At least 1 core clinical characteristic must be optic neuritis, acute myelitis with LETM\*, or area postrema syndrome
  - ☐ Dissemination in space (≥2 different core clinical characteristics)
  - ☐ Fulfillment of additional MRI requirements, as applicable (see chart below)

### **AND**

☐ Alternative diagnoses have been excluded (e.g., multiple sclerosis, sarcoidosis, cancer, chronic infection, etc.)

#### **AND**

- ☐ Member must have failed at least <u>ONE</u> of the following treatment options:
  - Oral therapy with azathioprine, methotrexate, mitoxantrone or mycophenolate
  - ☐ Member has required plasmapheresis or plasma exchange (PE) or intravenous Immunoglobulin (IVIG).

#### **Core Clinical Characteristics of NMOSD**

- Optic neuritis
- Acute myelitis
- Area postrema syndrome: episode of otherwise unexplained hiccups or nausea and vomiting
- Acute brainstem syndrome
- Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions
- Symptomatic cerebral syndrome with NMOSD-typical brain lesions

#### **Core Clinical Characteristics of NMOSD**

- Acute optic neuritis: requires brain MRI showing (a) normal findings or only nonspecific white matter lesions, OR (b) optic nerve MRI with T2-hyperintense lesion or T1-weighted gadolinium- enhancing lesion extending over >1/2 optic nerve length or involving optic chiasm
- Acute myelitis: requires associated intramedullary MRI lesion extending over ≥3 contiguous segments (LETM) OR ≥3 contiguous segments of focal spinal cord atrophy in patients with history compatible with acute myelitis
- Area postrema syndrome: requires associated dorsal medulla/area postrema lesions
- Acute brainstem syndrome: requires associated peri-ependymal brainstem lesions

(Continued on next page)

Medication being provided by: Please check applicable box below.
□ Location/site of drug administration:
NPI or DEA # of administering location:
<u>OR</u>
□ Specialty Pharmacy
For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.
**Use of samples to initiate therapy does not meet step edit/preauthorization criteria.**  *Previous therapies will be verified through pharmacy paid claims or submitted chart notes.