## SENTARA COMMUNITY PLAN (MEDICAID)

## MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

# SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (sJIA) OR ADULT ONSET STILL'S DISEASE (AOSD)

<u>Drug Requested</u>: Ilaris<sup>®</sup> (canakinumab) (J0638) (Medical) (Non-Preferred)

## **Medication can ONLY** be provided by a Physician's office

MEMBER & PRESCRIBER IN	NFORMATION: Authorization may be delayed if incomplete.				
Member Name:					
Member Sentara #:	Date of Birth:				
Prescriber Name:					
	Date:				
Office Contact Name:					
	Fax Number:				
DEA OR NPI #:					
DRUG INFORMATION: Author	orization may be delayed if incomplete.				
Drug Form/Strength:					
Dosing Schedule:	Length of Therapy:				
Diagnosis:	ICD Code, if applicable:				
Weight:	Date:				
☐ Standard Review. In checking this b	box, the timeframe does not jeopardize the life or health of the member aximum function and would not subject the member to severe pain.				

Recommended dosage: every 4 weeks SQ: 4mg/kg (with a maximum of 300mg) > 7.5kg

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial approval - 3 months.** For <u>continued 12-month approval</u>, please refax form with documentation of CRP or ESR along with progress notes to document therapy effective.

(Continued on next page)

□ D	OIAGNOSIS: Systemic Juvenile Idiopathic Arthritis (sJIA)
	Members must be aged 2 years - 17years
	AND
	Member must have had persistent sJIA activity for a minimum of six (6) months
	Date of diagnosis must be noted
	AND
	Member must have trial and failure of NSAIDs and corticosteroids for > 3 months consecutively within the last 4 months (paid claims will be reviewed for verification)
	AND
	Member must have had $\geq 5$ active joints with concomitant fever for at least 2 weeks within the last 3 months of this request
	<u>OR</u>
	Member must have had > 2 active joints with concomitant fever for at least 5 days and trial of prednisone or equivalent dosed at 0.5mg/kg/day or 30mg/day within the last 3 months of this request
	AND
	Member must have had CRP (>15 mg/L) within the last 2 months of this year
	AND
	Member must have had ESR (>45mm/hr) within the last 2 months of this year
	AND
	Member must have had fever > 380 C or 100.40 F for at least 2 weeks within the last 2 months of this request
	AND
	Member must have documented failure of Kineret <sup>®</sup> & either Arcalyst <sup>®</sup> or Actemra <sup>®</sup> (failure is defined as paid claims of Kineret <sup>®</sup> & Actemra <sup>®</sup> for at least 6 months AND lab values above did not respond to the preferred drug)
	<u>OR</u>
	Member has history of anaphylactic reaction to Kineret <sup>®</sup> , Arcalyst <sup>®</sup> or Actemra <sup>®</sup> [anaphylaxis is defined as an emergency department (ER/ED) visit due to throat or tongue swelling and/or shortness of breath] or development of skin reactions that lead to Stevens Johnson syndrome.

(Continued on next page)

□ D	OIAGNOSIS: Adult Onset Still's Disease (AOSD)
	Members must be aged $\geq 18$ years and $\leq 75$ years
	AND
	Member must meet two of the following:  □ Fever >39 °C, lasting 1 week or longer  □ Arthralgia or arthritis, lasting 2 weeks or longer  □ Typical rash  □ Leukocytosis >10,000/mm3 with >80% polymorphonuclear cells
	AND
	Disease activity based on DAS28 of ≥3.2 at screening
	<u>AND</u>
	Member must have had CRP (>15 mg/L) within the last 2 months of this year
	AND
	Member must have had ESR (>45mm/hr) within the last 2 months of this year
	AND
	At least 4 painful and 4 swollen joints at least 2 weeks within the last 3 months of this request
	AND
	Trial and failure with glucocorticoids, stable dose of $\leq$ 10 mg/day (prednisolone or equivalent) for at least 4 weeks AND NSAIDs, at least 4 weeks (at least 2 weeks within the last 3 months of this request)
	<u>AND</u>
	Trial and failure Kineret® and either Arcalyst® or Actemra® (progress notes must be submitted for true failure)
To su	uthorization Approval: 1 year. Check below all that apply. All criteria must be met for approval. apport each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be ded or request may be denied. Check box below for the Diagnosis that applies.
For s	JIA:
	Documentation decrease ESR <30.mg/L AND ESR<13mm/h
	AND
	Numbered of swollen joints have decreased

(Continued on next page)

For	•	$\boldsymbol{\cap}$	0	$\mathbf{r}$	
HAP	$\Lambda$				•
TOT		$\mathbf{v}$	V.	v	•

□ Documentation decrease ESR <30.mg/L AND ESR<13mm/h

### **AND**

□ Numbered of swollen joints have decreased

#### **AND**

□ DAS28 have decreased <2.6

Progress notes and labs documenting anaphylactic reaction or development of SJS must be submitted.

## Medication being provided by a Specialty Pharmacy - PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\*