

# SENTARA HEALTH PLANS

## MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-668-1550.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

**For Medicare Members:** Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Additional indications may be covered at the discretion of the health plan.

**Drug Requested:** Daxxify® (daxibotulinumtoxinA-lanm) (J0589) (Medical)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

• **Cosmetic indications are EXCLUDED**

**Recommended Dosing:** IM: Inject 125 to 250 units as a divided dose among affected muscles. Dose and number of injection sites should be individualized based on prior treatment, response, duration of effect, and adverse events. Dosage may be adjusted in 50- to 75-unit increments based on individual response; total recommended dose in a single treatment session: 125 to 250 units. Do not administer more frequently than every 3 months.

**Maximum Quantity Limits:** 250 units in a 3-month period

(Continued on next page)

