

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Cosentyx® SQ (secukinumab) (self-administered) (Pharmacy)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

**Cosentyx® is available under both Medical and Pharmacy benefits
(Please select correct PA form)**

| DIAGNOSIS | Recommended Dose |
|--|--|
| <input type="checkbox"/> Active Ankylosing Spondylitis | <ul style="list-style-type: none">Five syringes/pens in the initial 28 daysOne syringe/pen per 28 days after induction period. If patient continues to be symptomatic on above may increase to two syringes/pens per 28 days. |
| <input type="checkbox"/> Active Non-Radiographic Axial Spondyloarthritis (nr-axSpA) – with objective signs of inflammation | <ul style="list-style-type: none">Four syringes/pens in the initial 28 daysOne syringe/pen per 28 days after induction period or one syringe/pen per 28 days without induction |
| <input type="checkbox"/> Active Psoriatic Arthritis | <ul style="list-style-type: none">Five syringes/pens in the initial 28 daysOne syringe/pen per 28 days after induction period |

(Continued on next page)

| DIAGNOSIS | Recommended Dose |
|--|--|
| <input type="checkbox"/> Active Enthesitis-related arthritis (ERA) | <ul style="list-style-type: none"> Weeks 0, 1, 2, 3, and every 4 weeks thereafter. Weight \geq 15 kg and $<$ 50 kg the dose is 75 mg. Weight \geq 50 kg the dose is 150 mg |
| <input type="checkbox"/> Moderate to Severe Chronic Plaque Psoriasis | <ul style="list-style-type: none"> Ten syringes/pens in the initial 28 days Two syringes/pens per 28 days after induction period <p>Pediatric:</p> <ul style="list-style-type: none"> < 50kg: Five pediatric 75mg syringes in initial 28 days. One syringe per 28 days after induction period >50kg: Five pediatric 75mg syringes in initial 28 days. Two syringes per 28 days after induction period |
| <input type="checkbox"/> Moderate to Severe Hidradenitis Suppurative (HS) | <ul style="list-style-type: none"> 300 mg at weeks 0, 1, 2, 3, 4 and every 4 weeks thereafter. Dose may be increased to 300mg every 2 weeks if patient does not adequately respond |

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis: Active Ankylosing Spondylitis

- Member is 18 years of age or older
- Trial and failure of **BOTH** of the preferred drugs below (**check each tried**):

| | |
|---|----------------------------------|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd) | <input type="checkbox"/> Enbrel® |
|---|----------------------------------|

Diagnosis: Active Psoriatic Arthritis

- Member is 2 years of age or older
- Trial and failure of **TWO (2)** of the preferred drugs below (**check each tried**):

| | | |
|--|----------------------------------|---|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd) | <input type="checkbox"/> Enbrel® | <input type="checkbox"/> Pyzchiva® syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor) |
|--|----------------------------------|---|

(Continued on next page)

Diagnosis: Moderate to Severe Chronic Plaque Psoriasis

- Member is 6 years of age or older
- Member has a diagnosis of moderate to severe plaque psoriasis who is a candidate for systemic or phototherapy
- Member must have a previous failure on a topical psoriasis agent
- Trial and failure of **TWO (2)** of the preferred drugs below:

| | | |
|--|----------------------------------|---|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd) | <input type="checkbox"/> Enbrel® | <input type="checkbox"/> Pyzchiva® syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor) |
|--|----------------------------------|---|

Diagnosis: Non-Radiographic Axial Spondyloarthritis

- Member is 18 years of age or older
- Member has a diagnosis of Active Non-radiographic Axial Spondylarthritis (nr-axSpA) with objective signs of inflammation

Diagnosis: Active Enthesitis-related arthritis (ERA)

- Member is 4 years of age or older
- Member has a diagnosis of active Enthesitis-Related Arthritis
- Trial and failure of at least **two (2)** NSAIDs **OR**
- Use of NSAIDs is contraindicated in patient
- Trial and failure of **BOTH** of the preferred drugs below:

| | |
|--|----------------------------------|
| <input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd) | <input type="checkbox"/> Enbrel® |
|--|----------------------------------|

Diagnosis: Hidradenitis Suppurativa (HS)

- Member is at least 18 years old
- Member has a diagnosis of moderate to severe Hidradenitis Suppurativa
- Trial and failure of adalimumab-adbm (Boehringer Ingelheim) **OR** Hadlima® (adalimumab-bwwd)

Medication being provided by Specialty Pharmacy - PropriumRx

Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.