SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this</u> request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not</u> <u>complete, correct, or legible, the authorization process may be delayed.</u>

Drug Requested: (Select drug below)

deferiprone (Ferriprox [®]) tablets	□ Ferriprox [®] (deferiprone) solution	Ferriprox [®] 2-day (deferiprone) tablets	
MEMBER & PRESCRIBE	R INFORMATION: Authorizatio	n may be delayed if incomplete.	
Member Name:			
	Date of Birth:		
Prescriber Name:			
Prescriber Signature:		Date:	
Office Contact Name:			
Phone Number:	Fax Number:		
DEA OR NPI #:			
	uthorization may be delayed if incom		
Drug Form/Strength:			
Dosing Schedule:	Length of 7	Length of Therapy:	
Diagnosis:	ICD Code,	ICD Code, if applicable:	
Weight:	Date:	Date:	

Quantity Limits: Maximum 99 mg/kg/day (actual body weight) in two divided doses

CLINICAL CRITERIA: Check below all that apply. <u>All criteria must be met for approval</u>. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. Check the diagnosis below that applies.

Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome Initial Authorization: 6 months

- □ Member is 3 years of age or older
- □ Member has a diagnosis of transfusional hemosiderosis due to thalassemia syndrome (i.e., transfusion of \geq 100 mL/kg of packed red blood cells, approximately 20 units for a 40 kg patient)

- □ Member's serum ferritin levels are consistently >1,000 mcg/L (submit serum ferritin labs done within the last 30 days)
- □ Member's current weight: _
- □ Member is currently established on <u>ONE</u> of the following chelator therapies for at least 90 consecutive days (verified by pharmacy paid claims):
 - □ deferoxamine (Desferal)
 - □ deferasirox (Exjade, Jadenu)
- □ Member continues to have <u>ONE</u> of the following after three (3) months of chelator therapy:
 - □ Serum ferritin in excess of 2,500 mcg/L
 - □ Liver iron concentration is >7 mg Fe/g dry weight (submit liver biopsy, MRI or other FDAapproved test results)
 - □ Cardiac magnetic resonance imaging (MRI T2*) is ≤ 10 ms (submit MRI T2* lab results)
- □ Baseline absolute neutrophil count (ANC) is $> 1.5 \times 10^{9}$ /L and ANC will continue to be monitored weekly while on therapy (submit current labs)
- □ If requesting brand Ferriprox, documentation of trial and intolerable life-endangering adverse event with generic deferiprone must be submitted
- □ Ferriprox solution may be approved for members aged 3-10 years only. If requesting Ferriprox solution for members ≥ 11 years of age, documentation that member is unable to ingest any solid dosage form must be submitted

Diagnosis: Transfusional hemosiderosis due to thalassemia syndrome

Reauthorization: 12 months.

- □ Member's ANC is > 1.5×10^{9} /L (submit current lab results)
- □ Liver iron concentration is ≤ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test results)
- □ Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L
- □ Serum ferritin has decreased by $\ge 20\%$ from baseline or has been maintained at a level that is $\ge 20\%$ below baseline level (submit current serum ferritin labs)

Diagnosis: Transfusional iron overload in members with sickle cell disease or other anemias

Initial Authorization: 12 months

- □ Member is 3 years of age or older
- Member has a diagnosis of transfusional iron overload associated with sickle cell disease or other anemia diagnosis
- □ Baseline liver iron concentration > 7 mg of Fe/g dry weight (submit current MRI results)

(Continued on next page)

- □ Member has received no less than 20 transfusions of RBCs
- □ Baseline absolute neutrophil count (ANC) is > 1.5×10^{9} /L and ANC will continue to be monitored weekly while on therapy (submit current labs)

Diagnosis: Transfusional iron overload in members with sickle cell disease or other anemias

Reauthorization: 12 months.

- □ Liver iron concentration has decreased by at least 4 mg of Fe/g dry weight from baseline or has been maintained at a level that is at least 4 mg of Fe/g dry weight below baseline level since last approval (submit current MRI results)
- □ Member's ANC is > 1.5×10^{9} /L (submit current lab results)
- □ Treatment will be withheld if serum ferritin falls consistently below 500 mcg/L

Medication being provided by a Specialty Pharmacy - PropriumRx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.

<u>Previous therapies will be verified through pharmacy paid claims or submitted chart notes.</u>