

# SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** Sunosi® (solriamfetol)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

- Diagnosis of narcolepsy **must** be in accordance with the third edition of the International Classification of Sleep Disorders (ICSD-3), which is a fully revised version of the American Academy of Sleep Medicine's manual of sleep disorders nosology, published in cooperation with international sleep societies and is the key reference work for the diagnosis of sleep disorders.
- Sunosi® for narcolepsy will **NOT** be approved in conjunction with Lumryz™/Xyrem®/Xywav® or Wakix®. The Health Plan considers the use of concomitant therapy with Sunosi and Lumryz™/Xyrem®/Xywav® or Wakix® to be experimental and investigational. Safety and efficacy of these combinations has **NOT** been established and will **NOT** be permitted. In the event a member has an active Lumryz™/Xyrem®/Xywav® or Wakix® authorization on file, all subsequent requests for Sunosi® will **NOT** be approved.

**\*\*The maximum daily dose for this medication is 150 mg/day and requires renal dose adjustment\*\***

### Dosing Recommendations Based on Renal Function

Estimated GFR	Initial Dose	Maximum Dose
30-59 mL/min	37.5 mg once daily	75 mg once daily
15-29 mL/min	37.5 mg once daily	37.5 mg once daily
<15 mL/min	Not Recommended	Not Recommended

(Continued on next page)

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**DIAGNOSIS:** Please check **ONE** of the applicable diagnoses below

☐ **Narcolepsy** (All applicable boxes below must be met to qualify)

- ☐ Member 18 years of age or older
- ☐ Member does **NOT** have a history of alcohol, drug or stimulant abuse
- ☐ Members with a history of psychosis of bipolar disorders are being observed for possible emergence or exacerbation of psychiatric symptoms
- ☐ Member's blood pressure and heart rate has been assessed and is adequately controlled prior to initiating treatment and will be monitored regularly during treatment
- ☐ Member has a diagnosis of excessive daytime sleepiness associated with narcolepsy with or without cataplexy (**MSLT confirming diagnosis of narcolepsy must be submitted**)
- ☐ Member has failed a 30-day trial of modafinil or armodafinil (**verified by chart notes or paid pharmacy claims**)

☐ **Obstructive Sleep Apnea** (All applicable boxes below must be met to qualify)

- ☐ Member is 18 years of age or older
- ☐ Member does **NOT** have a history of alcohol, drug or stimulant abuse
- ☐ Members with a history of psychosis of bipolar disorders are being observed for possible emergence or exacerbation of psychiatric symptoms
- ☐ Member's blood pressure and heart rate has been assessed and is adequately controlled prior to initiating treatment and will be monitored regularly during treatment
- ☐ Member has a diagnosis of excessive daytime sleepiness associated with obstructive sleep apnea (**polysomnogram confirming diagnosis of OSA must be submitted with request**)
- ☐ Member has failed a 30-day trial of modafinil or armodafinil (**verified by chart notes or paid pharmacy claims**)
- ☐ Medication is **NOT** being used as primary treatment of underlying airway obstruction
- ☐ Standard treatment(s) for the underlying obstruction (e.g., with continuous positive airway pressure [CPAP], bi-level positive airway pressure [BiPAP]) have been used for one month or longer and have been properly titrated
- ☐ Member is fully compliant with ongoing treatments(s) for the underlying airway obstruction

**Medication being provided by Specialty Pharmacy – Proprium Rx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****