



GUIDE TO SUCCESSFUL BREASTFEEDING

sentara.com/maternity



TABLE OF CONTENTS

4	ADVANTAGES TO BREASTFEEDING For the Newborn For the Mother	16	SUPPORTING YOUR BREASTS Helping Your Baby Latch Signs Of An Effective Latch Signs Of An Ineffective Latch Nipple Pain
6	ANATOMY OF THE BREAST Milk Supply Colostrum: Baby's First Milk Factors That Affect Milk Supply Consequences Newborn Stomach Capacity	18	LENGTH AND FREQUENCY OF FEEDINGS Burping
10	BEGINNING TO BREASTFEED Baby-Led Attachment Skin to Skin Early Feeding Cues Crying To Awaken a Sleepy Baby	20	SIMPLE GUIDE BREASTFEEDING IN THE FIRST WEEK
14	POSITIONING AND LATCHING ON Baby-Led Football Hold Cross Cradle Hold Side-lying Cradle Hold	22	HOW TO TELL IF BABY IS GETTING ENOUGH Growth Spurts Manual Expression of Breast Milk Draining the Milk Reservoirs Assisting the Milk Ejection Reflex
		24	COMMON CONCERNS Nipple Soreness Flat/Inverted Nipples

	Treatment for Engorgement		Simple/Easy Breast Milk Storage Guidelines
	Breast/Nipple Swelling		Medically Indicated
	Reverse Pressure Softening		Reasons to Supplement
	Areolar Compression		Guidelines for Complementary Feedings
	Clogged Milk Ducts/Breast Infection		
	Treatment for Clogged Milk Ducts/	41	BLOOD SUGAR MONITORING
	Breast Infection		
	Mastitis (Breast Infection)		ALTERNATIVE METHODS OF SUPPLEMENTATION
	Yeast Infection (candida albicans)/Thrush	42	Pace Bottle Feeding
	Breast Surgeries		Spoon Feeding
	Postpartum Depression, Mood and Anxiety Disorders		Cup Feeding
	Nutrition		Syringe Feeds (Finger Feeding)
	Medications		Weaning
	Caffeine		
	Alcohol and Drugs		ADDITIONAL RESOURCES
	Contraception	46	
	Safe Sleeping and Breastfeeding		REFERENCES
	Pacifier Use	47	
35	SPECIAL CONSIDERATIONS	48	SPECIAL INSTRUCTIONS
	Twins/Multiples		
	Late Preterm Infants	49	BABY'S SECOND NIGHT
37	MILK COLLECTION AND STORAGE	50	FIRST TWO WEEKS: DAILY BREASTFEEDING LOG
	Pumping Guidelines		
	Pump Settings		
	Renting a Breast Pump		
	Tips for Increasing Milk Production when Pumping	54	NOTES



CONGRATULATIONS

THE ADVANTAGES TO BREASTFEEDING

Congratulations on your decision to breastfeed your newborn. While breastfeeding is the most natural way to feed your baby, it is also a learning experience for the two of you. This book has been prepared to give you the confidence and the information you need to successfully breastfeed your baby.

The best reason to breastfeed your newborn is the strong desire to do so. The first several weeks are a period of adjustment for you and your baby. As the weeks continue and a routine begins to develop, the rough edges begin to smooth. The enjoyment you feel in your relationship with your baby grows stronger as each day passes.

FOR THE NEWBORN:

- Due to breastmilk's protective mechanisms (antibodies), breastfed babies are less likely to develop ear infections, respiratory illnesses and gastrointestinal diseases.
- Breastmilk is easily digested. Breastfed babies are rarely constipated and have a lower incidence of stomach upsets and diarrhea.
- The incidence of allergies is decreased.
- Breastfeeding is excellent for jaw and tooth development.
- It's difficult to overfeed a breastfed baby, which may decrease the incidence of obesity.
- Evidence indicates that breastfed babies have lower cholesterol levels as adults, and decreased incidence of adult onset diabetes and childhood cancers.
- Recent studies show breastfed babies have higher IQ's.
- **BABIES LOVE TO BREASTFEED.**

FOR THE MOTHER:

- Encourages bonding between mother and baby.
- Helps the mother's uterus return faster to its pre-pregnant size.
- Breastfeeding full time burns up to 500 calories/day, aiding in mom's after delivery weight loss.
- Encourages rest periods.
- Is more cost-effective and saves time.
- Decreases the chance of developing ovarian and breast cancer, and osteoporosis. The longer a woman breastfeeds, the lower the risk.

ANATOMY OF THE BREAST


When your baby begins sucking, milk flows from the back of the breasts forward. The milk travels down the ducts behind the areola (the dark skin behind the nipple). Your baby's mouth and jaw movements will press the milk out of the ducts. These compressions allow the milk to flow out of the nipple openings into the baby's mouth.

MILK SUPPLY

A mother's milk supply depends on frequent, regular stimulation and emptying of the breasts. The law of supply and demand will continue as long as you:

- Breastfeed on demand or every 2 to 3 hours.
- If separated from the infant; breast massage, hand expression, and pumping within the first 6 hours will establish milk supply.

This means more breastfeeding equals more milk.



Anatomy of the breast.

COLOSTRUM: BABY'S FIRST MILK

Also known as liquid gold, due to its color and benefits for baby. ALL mothers have colostrum, a sticky yellow or clear liquid you may or may not have noticed before delivery. It can leak from your breasts as early as 5 months into your pregnancy.

Babies need colostrum because it:

- Contains antibodies that help protect them from illness.
- Builds the immune system.
- Has a laxative effect that helps clear the baby's digestive tract and helps the baby pass the meconium.
- Is high in protein and is easily digested.
- Is present and ready for baby at birth.
- Becomes **transitional milk** (a mixture of colostrum and mature milk) on day 2 to 5.
- Becomes **mature milk** between days 5 to 7.

FACTORS THAT CAN AFFECT AND/OR DELAY YOUR MATURE MILK SUPPLY:

- Baby not nursing well/sleepy baby.
- Supplemental formula feeding.
- High blood pressure and swelling.
- Gestational diabetes.
- Cesarean section.
- Excessive blood loss.
- History of breast surgery especially breast reduction.
- Shallow latch on.
- Pumping less than 8 to 10 times in 24 hours.
- Early pacifier use.
- Delayed breastfeeding or pumping if you are separated from your baby.

Your nurse or lactation consultant may recommend and assist you with using a hospital grade breast pump to stimulate your milk supply. Pumping in place of missed feedings can help your body produce more of the hormones that make milk.

CONSEQUENCES OF GIVING BOTTLES AND FORMULA BEFORE BREASTFEEDING WELL ESTABLISHED:

- The baby may develop a preference for firmer, faster flow bottle nipples.
- Sucking on a bottle nipple causes the baby's tongue goes back in mouth to control fast flow from bottle nipple. Which is the opposite of breastfeeding during which baby's tongue must go forward to cover the lower gum and suckle at the breast.
- Sucking on a bottle nipple causes the baby's tongue to go back in mouth to control the fast flow from the bottle nipple which is the opposite of breastfeeding. During breastfeeding, baby's tongue must go forward to cover the lower gum and suckle at the breast.
- Less stimulation of the breast equals less milk produced.
- Increase risk of severe engorgement of breast tissue once milk does come in.

LET DOWN REFLEX

Breastfeeding on demand or every 2 to 3 hours stimulates your body to release the two hormones that help your body produce milk for your baby.

- **Prolactin** increases the milk supply and helps with relaxation. It is also called the mothering hormone.
- **Oxytocin** contracts the muscle around your milk cell, releasing the milk into the ducts. Also known as the "milk ejection reflex" or "let down" reflex.

Some women feel a tingling sensation or an ache in their breast as the milk lets down, while others feel nothing. Once your milk is in, your breast may:

- Feel firm to hard and lumpy.
- Feel warm to hot.
- Feel uncomfortable to painful.
- Leak milk.

As you breastfeed your baby, your breast should soften and you should listen for swallows. Another effect of your breastfeeding hormones is an increase in the level of bonding and the peacefulness and comfort you feel as you breastfeed. Research shows that breastfeeding actually:

- Reduces stress hormones in both mom and baby.
- Helps both mom and baby rest better and go to sleep easily after breastfeeding.


NEWBORN STOMACH CAPACITY

Most full-term newborns:

- Have a strong sucking need.
- Are not hungry in the first day or two of life.
- Swallow amniotic fluid prior to birth.

Are born with extra fluid and fat stored to help prevent any dehydration until breastfeeding and breast milk gets established.

The following “belly ball” educational tool helps demonstrate a newborn’s stomach capacity.



The diagram illustrates the relative stomach capacity of a newborn at different stages of life using three objects: a small marble, a shooter marble, and a ping pong ball. It also includes a softball for comparison with an adult's stomach capacity. A logo for Sentara Women's Health Source is present.

Babies Were Born To Be Breastfed!	
Sentara Women's Health Source	
1 Small Marble	= Stomach capacity of a newborn on Day 1 5-7 mL
Shooter Marble	= Stomach capacity on Day 3 22-27 mL
Ping Pong Ball	= Stomach capacity on Day 10 60-81 mL
Softball	= Stomach capacity of an adult

- Ask that your baby’s bath be delayed until you can at least attempt to breastfeed, even if it is in the recovery room after a Cesarean birth.
- Please avoid bottles for the first feeding, even if supplementation is medically necessary. Introducing an artificial nipple before breastfeeding is established can make it more difficult for the baby to learn how to latch on and suck properly on the breast. Your nurse can assist with alternative feeding methods such as a cup or syringe feed (See page 42-43).



BEGINNING TO BREASTFEED

BABY-LED ATTACHMENT

All newborns are “hard-wired” to breastfeed and possess the instincts needed to latch-on to mother’s breast.

Breastfeeding happens more naturally when both mom and baby are calm and comfortable. This can be accomplished easily by placing baby, wearing only his diaper, on moms bare chest, “skin to skin” or “heart to heart.” Baby feels his mother’s breathing and heartbeat, just like when he was still attached in the womb. Mom can snuggle her baby and respond to his early feeding cues.

SKIN TO SKIN AS SOON AS POSSIBLE AFTER BIRTH HAS THESE POSITIVE EFFECTS:

- Awakens baby’s rooting instincts. Babies “root” for the breast by opening mouth wide and bobbing head or squirming towards nipple.
- Many babies will self attach to the nipple with a wide open mouth when rooting.
- Baby is less likely to cry and is more calm and content.
- Baby maintains his body temperature, has normal respiratory and heart rates and stable blood sugars.

- When separated from their mother, babies will cry more, have lower blood sugars and take longer to adjust to their new environment.
- Studies show that skin to skin contact for at least 20 minutes after birth increases the duration that babies are exclusively breastfed.

EARLY FEEDING CUES

Mother needs to learn to feed in response to her baby's early feeding cues such as:

- Moving head side to side with a wide open mouth.
- Licking and sucking motions with tongue and mouth.
- Trying to get hands or fingers in mouth.

CRYING

Crying is a late feeding cue, and a crying baby has a harder time trying to latch on to your breast and have an organized suck. A crying baby also can wear himself out and go to sleep quickly once brought to your breast.






A good way to calm a crying baby is to let the baby suck on your clean finger. Just gently place your finger in his mouth and with the pad of your finger gently massage the roof of his mouth which will elicit his sucking reflex. Gently rocking baby and making shushing sounds as loud as his cries also helps calm baby.

TO AWAKEN A SLEEPY BABY

Most newborns are typically very sleepy the first 24 hours after birth. Their journey into the world often just wears them out. They are overwhelmed by so many new sensations, touch, sights, sounds, bathing, etc. When they go to your breast, they are comforted by your smell, breathing and heartbeat and would rather sleep than eat. Let baby have some snuggle time skin to skin, and that often will wake up their instinct to breastfeed. When it is time to wake a sleepy baby for a feeding, here are some tips:

- Unwrap the blanket, remove hat, and change the diaper if necessary.
- Dim the light, so the baby will open his or her eyes.
- Talk to your baby.
- Play with the baby's mouth or lips.
- Exercise baby's arms and legs, or do gentle "baby sit-ups".
- Gently rub baby's back.
- Wipe baby's face with a cool cloth.
- Undress the baby down to the diaper.



POSITIONING AND LATCHING ON

- Bring the baby to breast level using pillows or blankets.
- Baby should be facing Mom (tummy to tummy).
- Do not lean over baby, remember baby to breast, not breast to baby.
- Baby's nose to mom's nipple.
- Baby led attachment, don't force nipple into mouth but be patient and wait for rooting and latching on with wide open mouth.
- Support your breast throughout the feeding.
- Once latched the infant's mouth should be wide open, lips flared out, cheeks rounded and tongue down over the gum line.
- Mom should feel strong tugging, but no biting or pinching pain throughout the feeding.

POSITIONS AT THE BREAST

There are several different positions for breast feeding. Experiment to find what is comfortable and what works best for you and your baby. It is a good idea to rotate these positions to help prevent sore nipples. For all positions you can use pillows or a boppy to raise baby to breast level. This will allow you to more fully relax your arms and shoulders after the baby has latched on. Remember leaning forward over your baby inhibits his rooting reflexes and will leave you with an aching back.

BABY-LED

Recline in the bed or on the sofa, about half way back. Place your baby skin to skin with you, between your breasts. Your baby will begin to root and move to the breast. You provide support as baby finds the breast and latches by himself.



FOOTBALL HOLD:

This works best with Mom in a comfortable chair. It is especially helpful for women who have had a cesarean, for those who have large breasts or for women with a small baby. Make sure your hand is between the baby's shoulder blades, with your fingers supporting his head.



CROSS CRADLE HOLD:

The baby's body is supported by mom's forearm and baby's head is supported by mom's hand. Mom's arm that goes across her body supports the baby and mom's other hand supports her breast.



SIDE-LYING:

This encourages Mom to rest during feeds. It works well for women with large breasts or women who are uncomfortable sitting after delivery.



CRADLE HOLD:


Position baby's head on your forearm just below the crook of the elbow. Your hand supports baby's bottom with baby's stomach turned towards your stomach. This keeps baby's body in a straight line. Your opposite hand supports the breast in a "C" hold.



SUPPORTING YOUR BREASTS

You should support your breasts during your baby's feedings for the first few weeks. Your newborn's suck is not strong enough to support the weight of your breast at first. When you hold your breasts, you can use either a "C" or a "U" hold. Be sure your fingers are far enough away from the areola; so they don't interfere with the ability of your baby's gums to compress the milk ducts that are behind your areola.

Although your baby's nose may be touching your breast, all newborns have flared nostrils, so he can still breathe if his nose is on your breast. If he can't breathe, he can move his head. If a baby is sucking, he is breathing. It is not necessary to push firmly down on your breast tissue to clear the space at the nose. This often adds tension to your nipple, increasing nipple soreness. Many very large breasted women do need to be sure the baby's nose is not covered too much. A good tip for large breasted women is to place a rolled washcloth or receiving blanket under your breast. This supports the breast and lifts the nipple up, making it easier to latch your baby on.



HELPING YOUR BABY LATCH

- Hold your baby close to your body with his tummy facing yours and his nose across from your nipple.
- Gently tickle his upper lip with the tip of your nipple and WAIT for a wide open mouth like a yawn.
- Quickly pull him in close, chin first, so he gets a mouthful of breast.

Do not try to force your nipple into his mouth.

- This will cause baby to clamp down on your nipple leading to a painful latch and a decreased flow of colostrum/breast milk.
- A shallow latch also leads to pinched, cracked, and very painful nipples.



HELPING YOUR BABY LATCH CONTINUED

Remember – baby to breast, not breast to baby.

Leaning over your baby inhibits his rooting instincts. Leaning back and bringing baby onto your breast helps awaken his rooting instincts.

SIGNS OF AN EFFECTIVE LATCH:

- Initial mild discomfort on the nipple quickly subsides.
- You feel a rhythmic tug on your nipple as baby suckles.
- Baby's lips are rolled outward.
- Baby's nose, chin, lower lip and possibly cheeks are touching the breast.
- Baby's jaw and ear moves as they suck.
- You see and hear (once milk is in) swallowing.
- Your nipple appears rounded, elongated after the feeding.

SIGNS OF AN INEFFECTIVE LATCH:

- Baby's mouth is only on nipple and small amount of areola.
- You experience discomfort throughout the feeding.
- As the baby nurses you hear clicking, smacking sounds.
- After breastfeeding your nipple appears pinched or misshaped.
- You experience prolonged nipple soreness and/or trauma.

NIPPLE PAIN

When your baby first latches on, you may feel some tenderness due to nipple stretching; however, this should go away within 30-40 seconds. Painful sucking usually indicates a shallow latch. You should re-latch, but be careful not to just pull baby off. It is best to break the suction first by inserting your finger in the side of the baby's mouth, then remove him from your breast.

Persistent nipple pain that is uncorrected by a deeper better latch could indicate an issue with the baby's oral anatomy. Some baby's have a lack of tongue mobility due to a tight frenulum (the tissue that connects the baby's tongue to his base of his mouth) also called "tongue tie." Your baby's Pediatrician will evaluate if any intervention is necessary to correct this.

LENGTH AND FREQUENCY OF FEEDINGS

Babies need 8 to 12 feedings every 24 hours in the first 4 to 6 weeks.

- Time a feeding from the beginning of one feeding to the beginning of the next.
- If your baby has gone more than 3 hours without a feeding, **wake your baby.**
- During the first few days of life, the length and frequency of your baby's feedings may vary greatly. At times your baby may only nurse a few minutes and other times he/she may nurse longer.
- A feeding session can take 40 minutes or more in the first weeks while babies are still sleepy.
- Your baby may also have periods of **cluster feedings** when he/she is more awake and wanting to nurse much more frequently. This often occurs in the evening and or night. Remember to watch your baby, not the clock and breastfeed according to his feeding cues.

Offer both breasts at each feeding. Alternate which breast you begin each feeding with.

- Some babies will consistently nurse both breasts, while others will not.
- Consider the first breast the “meal” and the second breast the “dessert.”
- Babies nurse most vigorously at the beginning of a feeding, then slower at the end.
- There is more fat content at the end of each feeding. Switching breast too soon can possibly not satisfy baby’s hunger and lead to more fussiness and more frequent feedings.
- If your baby only nurses from one breast and falls asleep, with the next feeding, start on the other breast.

Limiting breastfeeding times will delay your milk coming in, cause engorgement or decrease your supply.

- Only your baby can tell you when he/she is hungry or full. Watch your baby and not the clock to assess when the feeding is done. Once satisfied, most babies will come off the breast or fall asleep.
- Limiting feedings in the first few days will delay the transition from colostrum to milk, as well as limit the amount of colostrum your newborn will receive.
- Longer feedings will increase your prolactin level, enabling you to make more milk for the next feeding, as well as emptying your breasts (remember supply and demand).
- If breastfeeding hurts, call for help (see back of booklet).

BURPING

- After feeding on one breast, attempt to burp your baby over your shoulder, lying on your lap or sitting up in your lap.
- Gently pat or rub your baby’s back for several minutes.
- If your baby burps, air that leaves the stomach will make room for more. It also helps to renew your baby’s interest in feeding from the second breast.
- If your baby does not burp and is awake, don’t worry. Go ahead and offer the second breast.
- At the end of the feeding, you may try to burp your baby if he or she is awake. If your baby is asleep, do not awaken to burp; just lay your baby on its back as recommended by the American Academy of Pediatrics.

A SIMPLE GUIDE TO BREASTFEEDING IN THE FIRST WEEK

	DAY 1	DAY 2	DAY 3
BREAST MILK	A few drops to a teaspoon per feeding, gold, clear	A few drops to a teaspoon per feeding, gold, clear	More colostrum with some milk, lighter golden color more creamy
BREASTS	Soft	Soft	Heaviness, soft with areas of fullness before feedings
BABY BEHAVIOR	Sleepy, multiple attempts, 1 or more successful feeds	More alert, 8-12 attempts, several successful feeds, cluster feeds	Alert for feedings, 8-12 feeds, fussy at night
SUCKLING AND SWALLOWING	Bursts of sucks followed by pause	Bursts of sucks followed by pause	Bursts of sucks, followed by pause, swallowing every 1-3 sucks
LENGTH OF FEED	5-45 minutes each side	5-45 minutes each side	5-45 minutes each side
URINE	1 wet diaper	2 wet diapers	3 wet diapers
STOOL	Minimum of one but usually several dark black, sticky meconium stools	Minimum of two but usually several dark black, sticky meconium stools	Usually 1-2 or more dark green or slightly brownish stools
WEIGHT	Weight Loss	Weight Loss	Less Weight Loss

DAY 4	DAY 5	DAY 6	DAY 7
More colostrum with some milk, lighter golden color more creamy	Good volume of transitional milk, more opaque, light yellow and creamy	Good volume of transitional milk, more opaque, light yellow and creamy	Good volume of transitional milk, more opaque, light yellow and creamy
Heaviness, fullness throughout before feeding	Fullness throughout before feeding, softer after	Fullness throughout before feeding, softer after	Fullness throughout before feeding, softer after
Alert for feedings, 8-12 feeds, more satisfied after feeds	Alert for feedings, 8-12 successful feeds	Alert for feedings, 8-12 successful feeds	Alert for feedings, 8-12 successful feeds
Bursts of sucks, followed by pause, swallowing every 1-3 sucks	Bursts of sucks, followed by pause, frequent swallowing throughout feeding	Bursts of sucks, followed by pause, frequent swallowing throughout feeding	Bursts of sucks, followed by pause, frequent swallowing throughout feeding
10-30 minutes each side	10-30 minutes each side	10-30 minutes each side	10-30 minutes each side
4 wet diapers	5 wet diapers	6 wet diapers	6 wet diapers
Usually 1-2 or more slightly brownish or bright green loose stools	Usually several bright yellow, loose, maybe "seedy" stools	Several bright yellow, loose, maybe "seedy" stools	Several bright yellow, loose, "seedy" stools
Weight Stable	Weight Stable	Weight Gain	Weight Gain

HOW TO TELL IF BABY IS GETTING ENOUGH

- The baby nurses at least 8 times in 24 hours. (First day MAY be sleepy, recovering from birth.)
- The baby has as many wet diapers as days old. After 5 days, the baby has 6 very wet diapers every 24 hours. (See Breastfeeding Log, in the back of book for days 1-5). It may be difficult to assess wetness with some brands of disposable diapers. If unsure, place a folded tissue in the fresh diaper.
- Remember the first day an infant's stool is black and tarry. More brown green on day 2-3, becoming more yellow and seedy by day 5. It is even considered normal for your baby to stool after each feeding. Occasionally an older breastfed baby will have a stool every 3 to 5 days. As long as the stool is soft, this is normal.
- You can hear your baby swallowing (after your milk is in at 3 to 5 days) at a rate of one suck per second for the first half of the feed on that breast.
- If your baby is not eating well or is hard to awaken, call your pediatrician.

GROWTH SPURTS

- When a baby goes through a Growth Spurt, he or she may want to nurse more frequently.
- These occur approximately when baby is:
 - 2-3 weeks
 - 6 weeks
 - 3 months
 - 6 months old

Continue nursing as needed by your baby. Your body will begin to produce more milk to meet his/her nutritional needs within 24 to 48 hours.




MANUAL EXPRESSION OF BREAST MILK

DRAINING THE MILK RESERVOIRS

1. Position the thumb (above the nipple) and first two fingers (below the nipple) about 1" to 1-1/2" from the nipple, though not necessarily at the outer edges of the areola. Use this measurement as a guide, since breasts and areolas vary in size from one woman

to another. Be sure the hand forms the letter "C" and the finger pads are at 6 and 12 o'clock in line with the nipple. Note the fingers are positioned so that the milk reservoirs lie beneath them. Avoid cupping the breast.

2. Push straight into the chest wall. Avoid spreading the fingers apart. For large breasts, first lift and then push into the chest wall.
3. Roll thumb forward as if making a thumbprint and change finger pressure from middle to first finger at the same time. This rolling motion compresses and empties milk ducts without injuring sensitive breast tissue. Note the moving position of the thumb and fingernails as shown in the illustration.
4. Repeat rhythmically to completely drain milk ducts: position, push, roll... position, push, roll....




The Marmet Technique, Copyright 1978, revised 1979, 1981 and 1988.

Used with permission of Chele Marmet and The Lactation Institute, 16430 Ventura Blvd., Suite 303, Encino, California 91436 • Phone 818-995-1913.


AVOID THESE MOTIONS

1. Do not squeeze the breast, as this can cause bruising.
2. Avoid sliding hands over the breast as it may cause painful skin burns.
3. Avoid pulling the nipple and breast, which may result in tissue damage.



ASSISTING THE MILK EJECTION REFLEX: GENTLE MASSAGE

1. Massage the milk producing cells and ducts by pressing the breast firmly with the flat of the fingers into the chest wall, beginning at the top. Move fingers in a circular motion, concentrating on one spot at a time for a few seconds before moving on to another spot. Spiral around the breast toward the areola as you massage. The motion is similar to that used in a breast examination.
2. Stroke the breast area from the top of the breast to the nipple, using a light tickle touch. Continue the stroking motion to help you relax, which in turn will stimulate the milk ejection reflex.
3. Shake the breast while leaning forward so that gravity will help the eject.



COMMON CONCERNS

NIPPLE SORENESS

- Nipple soreness can often be avoided by proper positioning, latching on, and rotation of feeding positions. Baby's nose should be near the breast, and chin and cheeks should touch the breast. Baby's mouth should be opened wide, and lips flared outward.
 - To break the baby's suction, place your little finger between the gums before pulling baby off or pull down on the chin.
 - If your nipples become sore or cracked, feed on the least sore side first.
 - Avoid drying agents or blowing air on nipples. Apply a thin layer of lanolin to nipples after each feeding to soothe sore or cracked nipples, or hydrogels may be used.
 - Breast shells may be used if nipples are very tender and any contact with nipples causes greater discomfort.
 - Your nipples need only be washed with water. Rinsing during your daily shower is sufficient. If there are open cracks, nipples should be gently washed with a mild soap and rinsed with clean water daily.
 - You may find that feeding more often will help nipple soreness. Your baby may not nurse as vigorously if fed more often.
 - If your nipple sticks to the nursing pad, moisten the pad until it comes off easily. Change wet nursing pads with every feeding.
 - For persistent sore or cracked nipples, contact a lactation consultant.
- 24 Additional treatment may be necessary.

FLAT/INVERTED NIPPLES

Every woman's nipples are different. Some protrude (stick out), some are flat, and some are inverted (push in). All of these are normal. Remember to support/compress your breast tissue (sandwiching of breast) and bring baby to breast, not breast to baby. Most newborns will be able to latch regardless of the shape of the nipple when self-attaching and laying skin to skin with mom in a laid-back position.

Some moms and babies will need help with latching when they have flat or inverted nipples. Here are some helpful tips:

- Before feeding, gently tug and roll your nipple at its base to help draw your nipple out .
- Pumping your breasts using a pump prior to feeding can also help draw your nipple out.
- If your nipples are swollen see the section on reverse pressure softening pg. 26.
- Talk with your Lactation Consultant for latching aids if needed.

TREATMENT FOR ENGORGEEMENT

- **It is normal for your breasts to become warm, full, and heavy when your milk comes in. You may feel uncomfortable.**
- It is not normal for your breasts to feel red, hot and painful.
- **The best treatment for engorgement is prevention by frequent, on demand feedings and not setting time limits on feeding length (average newborns can empty the breast in 15 – 20 minutes).**
- If your nipples flatten out as your breasts become hard, your baby may have difficulty latching on. **You may need to release some milk from the ducts by massaging your breasts using warm, moist heat. Such as in the shower or with a compress, for only 2 to 3 minutes, or by expressing some milk by hand.**
- Gentle massage and compression during the breastfeeding session can help the infant remove more milk from your breast.
- Use deep breathing, soft music, or other techniques to relax before and during nursing.
- Most women are more comfortable with a bra on at this time. Do not wear tight, restrictive bras or clothing.
- **Ice packs after feedings are comforting (can use bags of frozen vegetables).**

If your breasts become severely engorged (skin tight, shiny, hard), you should call your Lactation Consultant for additional assistance. You may need a hospital grade/rental electric pump to help relieve this.


BREAST/NIPPLE SWELLING

- Often caused by the large volumes of the IV fluid some moms, with longer labors and epidurals and/or C-sections, receive.
- May interfere with baby's ability to latch.

Reverse pressure softening or areolar compression can be used to help relieve this swelling (used with permission from Jane Cotterman IBCLC).

REVERSE PRESSURE SOFTENING TWO THUMB METHOD

- Position the base of the thumbnail at the base of the nipple on opposite sides.
- Apply gentle inward pressure for 1-3 minutes until pits are formed.
- Change position as needed until the central areola is soft enough to try latching infant.



AREOLAR COMPRESSION

- Using 2 or 3 straight fingers on each side.
- Hold until edema is felt to give way.
- Observe for "pitting" representing edema.
- Apply pressure again behind the pitted spot.
- Continue to press and hold in this fashion working around the areola/ breast.
- Continue until areola is softened enough for infant to latch.



CLOGGED MILK DUCTS/BREAST INFECTION

- **Lumps in the breast may be a sign of clogged milk ducts.** This can occur when feedings are spread out (more than 4 hours in between feeds), a feeding is skipped, a breast is not emptied, or you are wearing restrictive clothing or an under wire bra.

TREATMENT FOR CLOGGED MILK DUCTS/BREAST INFECTION

- **If you feel a lump, apply warm, moist heat to the area for 5 to 10 minutes before feeding the baby and massage the area to assist in releasing milk.** Start on this affected side. Feed a minimum of 10 minutes on this side, then check after the feeding to see if the lump has disappeared.
- Position your baby so that the chin or nose is in the direction of the lump. This will be the strongest point of suction and will aid in releasing the milk.
- Feed frequently without time restrictions. If you must skip a feeding or if the baby extends its nap time, express some milk with a pump or by hand.

MASTITIS (BREAST INFECTION)

If a milk duct remains clogged after several feedings and the lump does not disappear, mastitis (a breast infection) may occur. Breast infections are more common in women who are fatigued, stressed, or not eating well. The symptoms may include:

- **Flu-like symptoms** such as general body aches and a headache, often with a **temperature of 101° or higher**. Chills often accompany the fever.
- There may be a **painful or tender reddened area on the breast**.
- It usually occurs in one breast only.
- If you have any of these symptoms call your obstetrician. You probably will be placed on antibiotics for 7 to 10 days. During this time it is important that you:
 - **Continue nursing** to keep the ducts empty (your milk is not infected).
 - **Get plenty of rest.**
 - **Drink plenty of fluids.**

YEAST INFECTION (CANDIDA ALBICANS)/THRUSH

A yeast infection can occur on the nipple if it is spread from your baby's mouth, after antibiotic therapy, or from damaged nipples left untreated. The symptoms include:

- Reddened, swollen, tender, or cracked nipples.
- Burning, itching, shooting pain deep in the breast, tenderness during or immediately after feedings.

If you have any of these symptoms call your obstetrician or your baby's pediatrician for antifungal treatment. It is important that you:

- Wash your hands before feeding your baby.
- Change your breast pads at each feeding.
- Boil items such as pacifiers, bottle nipples, breast shells, breast shields for breast pumps, and anything else that comes in contact with the baby's mouth and/or your milk for 5 minutes daily.
- **Decrease concentrated sweets in your diet. Add acidophilus to your diet (found in active culture yogurt or acidophilus milk).**
- **Treat you and your baby at the same time.**
- **Avoid freezing any milk pumped during a yeast infection as freezing does not kill yeast.**
- **Keep breastfeeding.**

BREAST SURGERIES

Please let your nurse and or lactation consultant know if you have had any breast surgeries. Breast implants or biopsies usually do not have any effect on milk production. However, breast lumpectomies or breast reduction surgeries often can affect milk production. The site www.bfar.org (breastfeeding after reduction) is a website with more information.

POSTPARTUM DEPRESSION, MOOD AND ANXIETY DISORDERS

Postpartum depression/anxiety is the #1 complication of pregnancy and childbirth. One in five new mothers will have this challenge. New breastfeeding moms can often have excessive anxiety and stress if breastfeeding's not going well and are often sleep deprived. Be sure you are getting enough rest, napping when your baby naps. If treatment is needed:

- Contact your doctor or obstetrician for recommendations for support groups or counseling.
- Medications are a safe treatment option with breastfeeding. Your doctor will know which medication will work best for you when breastfeeding.
- Studies show that moms who resolve breastfeeding issues, or are successfully breastfeeding right from the start, have less incidence of postpartum depression (postpartumva.org).

NUTRITION

- Continue with prenatal vitamins.

There is no special diet for breastfeeding mothers. **Most nursing moms are thirsty, drinking approximately 2 quarts of liquid per 24 hours.** If your urine is concentrated (dark amber with a strong, noticeable odor) you are not drinking enough!

Breastfeeding requires approximately 300 to 500 extra calories per day over your pre-pregnancy diet. Continue eating a regular wholesome diet, adding an extra serving of protein and calcium-rich foods. **There are no foods that you must eat and no foods that you cannot eat. Babies tend to handle mom's regular diet without trouble.** Do not go on a diet while lactating. Breastfeeding mothers may include fish, sushi and shellfish in their diets. The FDA advises eating fish known to be low in mercury.

Occasionally your baby may experience extreme gassiness/fussiness. Most foods pass through breast milk within the first three hours up to 24 hours. If you suspect a certain food in your diet is causing a problem, contact your Pediatrician or your lactation consultant for guidance. Some foods that you may suspect are:

- Dairy products (for example milk, cheese, yogurt).
- Green leafy vegetables.
- Overly spiced foods.
- Caffeinated beverage and chocolate. Keep in mind that caffeine levels peak in 45 minutes to 2 hours after ingestion.

Gassiness may also be a sign of thrush - a yeast infection in your baby's digestive system.

MEDICATIONS

Most medications are safe to take while breastfeeding. Consult with your Pediatrician or Lactation Consultant before taking a prescription medication while breastfeeding. Over-the-counter medications that are safe to take while breastfeeding include:

- Acetaminophen
- Ibuprofen
- Prenatal vitamins
- Stool softeners
- Robitussin

Cold medicines containing pseudoephedrine or antihistamines may decrease your milk supply.

Website: www.nlm.nih.gov/lactmed

CAFFEINE

Caffeine may keep your baby awake or make them fussy. Limit coffee to one or two cups a day and remember that sodas, teas, energy drinks and chocolate contain caffeine too.

ALCOHOL AND DRUGS

- The use of any illegal drugs while breastfeeding is contraindicated and could result in serious problems to your baby, not to mention legal consequences for you.
- Nicotine is considered a drug and having a baby is a great time to quit smoking. However, if you cannot stop smoking, the benefits to your baby outweigh any risks with breastfeeding. Formula fed babies of smoking mothers are much more likely to be sick, develop asthma and respiratory problems compared to breastfed babies of mothers who smoke. Please limit smoking to after a feed, outside and away from the baby and use a cover jacket to decrease the risk of second hand smoke, which increases the risk for SIDS.
- As with most everything, moderation is the key when drinking alcohol. The amount of alcohol transferred into breast milk is generally low. An adult will break down 1 ounce of alcohol in about 3 hours, so mothers who drink small amounts can return to breastfeeding/pumping when they are no longer feeling the effects. (Chronic heavy drinkers shouldn't breastfeed.)
- Rule of thumb: You may have one 12 ounce beer, a 6 ounce glass of wine or one ounce of alcohol. If drinking more than one serving, it is safest to "pump and dump" the next feeding.
- Alcohol does inhibit the release of oxytocin which will reduce letdown. This appears to be dose related and usually is seen with moderate to heavy drinking. Other studies show that alcohol changes the taste of milk and baby will nurse less and then nurse more as the alcohol level decreases.

CONTRACEPTION

During breastfeeding, the chance of getting pregnant may be lower; however, you can still get pregnant. It is a good idea to discuss contraception with your physician before or just after you give birth. Be aware of the many choices you have.

- Nonhormonal methods of birth control such as condoms, spermicides, diaphragms, IUD's, and sponges have the least effect on breastfeeding as there is nothing to pass into the milk supply, affect milk production or alter milk composition.
- Progesterone only birth control is the next choice as combination birth control with estrogen has been found to decrease milk production and is not recommended for use until after the baby is 6 months old.
- There is controversy over when to begin progesterone only birth control. There is a possible risk of decreased milk supply when hormonal contraception is started before milk production is well established. Most professional organizations and birth control drug manufacturers recommend waiting for at least 2 weeks postpartum to avoid a possible negative impact on lactation.
- Talk with your OB or Lactation Consultant regarding birth control options and how it could affect your milk production.

SAFE SLEEPING AND BREASTFEEDING

The American Academy of Pediatrics recommends that mothers and infants sleep near one another but in separate beds for the first 6 months of life and ideally, the first year of life. Room-sharing and breastfeeding decrease your infant's risk of SIDS.

- Studies show that when mothers and babies sleep in the same room, with baby in a bedside crib or bassinet, nighttime feedings are easier, mothers and babies get more sleep, and babies have a lower risk of SIDS.
- As a breastfeeding mother, you may be breastfeeding in bed. If there is a possibility of you falling asleep while breastfeeding, feed your baby in your bed rather than moving to a chair or sofa. The bed should have no pillows, blankets, sheets, people or other items that could block your baby's ability to breathe or cause overheating.
- There have been cases of babies suffocating or falling off unsafe places, like chairs, sofas, recliners or water beds.

Babies have been found suffocated when a parent in the bed has taken medications or is intoxicated and rolled over on top of their baby.

- Avoid baby's exposure to cigarette or cigar smoke, alcohol or illegal drugs.
- **Never leave your baby alone on an adult bed.**

Remember, baby sleeps Alone, on his/her Back, in the Crib.

PACIFIER USE

Studies have shown that early pacifier use can interfere with your infant's latch and/or your milk supply. It may seem like all you are doing is feeding your baby. Those first several weeks are very busy. Not only is your baby nursing for nutrition, he/she is establishing your milk supply for months to come. A pacifier should not be used to replace or delay a meal. It may be tempting to give a pacifier when it is convenient for your schedule. It is best to watch for hunger cues and feed on demand. Settle into an effective nursing routine. Your baby nurses differently at the breast than with a pacifier. At the breast, your baby takes the nipple into the junction of the soft and hard palate. A pacifier encourages thrusting motion with the tongue. Four weeks is an appropriate time to introduce a pacifier if you desire. Strive for success at the breast.

Infants are learning to breastfeed and need to have time to learn your breast. Your breasts need frequent stimulation to get your supply well established. Your milk "comes in" by day 5, but your supply may take up to 4 weeks to be well established.

Sucking on a pacifier releases the same digestive enzyme as sucking at the breast. This enzyme can cause the infant to feel like he/she is full and content. This can cause your infant to miss a feeding, and your breast will miss being stimulated.

The AAP recommendations for SIDS prevention suggests waiting until your milk supply and breastfeeding is well established before introducing a pacifier. Generally that could take up to 4 weeks.



SPECIAL CONSIDERATIONS

There are some circumstances that can take a mother and her newborn off the usual course for establishing breastfeeding such as:

- Prematurity, illness, or birth defect that results in admission to Neonatal Intensive Care Unit (NICU) or Special Care Nursery (SCN)
- Conditions that require blood sugar testing prior to feedings for infants that are still able to be in normal nursery or mother-baby rooming in.
- Newborn jaundice requiring phototherapy/bili lights.
- Infants who are unable to adequately breastfeed results in excessive weight loss or inadequate output.
- Inadequate milk supply due to maternal conditions such as breast surgeries, high blood pressure, excessive swelling, blood loss and recovery from Cesarean section or other surgery.
- Infants having withdrawal symptoms from maternal medications or drug/ nicotine use.

Many of these situations may require intravenous (IV) fluids, feeding of formula or pumped breast milk or medications.



TWINS/MULTIPLES

Many mothers delivering twins or even triplets are overwhelmed with the thought of how to manage breastfeeding. They often have babies that are premature or late preterm. They have to use a breast pump to help establish their milk supply. When the babies are unable to breastfeed due to their prematurity, low birth weight or decreased stamina, mothers often are encouraged to pump and supplement with their own expressed breastmilk or additional formula. Refer to section on “Alternate Methods of Supplementation” for different feeding methods.



The good news is there is no expiration date on babies being able to learn how to breastfeed, even after weeks of having tube feedings or bottles. Your babies’ nurses and the lactation consultants will be able to help you. Some important points to remember:

- Your body knows there is more than one baby to feed, and many moms with multiples can produce all the milk their babies need. (Just know it often takes a week of exclusively pumping to get that milk supply established.)
- Each baby has to learn individually how to breastfeed before trying to feed two babies at the same time.
- Working with a Lactation Consultant can help you develop a plan.

LATE PRETERM INFANTS

Babies born a few weeks early, 34-37 weeks gestation, are known as “late preterm infants.” They often can be in the normal newborn nursery but cannot be treated the same as a full term newborn. Because they are early, they are often small and immature in many of their abilities. Late preterm infants can have the following issues:

- Inability to maintain their body temperatures (because they have less fat deposits).

- Inability to have stable blood sugars (because they are burning more calories to stay warm).
- Are very sleepy, tire more easily because they do not have the same strength and stamina as a bigger more mature newborn.
- Can have more respiratory problems.
- Higher risk for jaundice.
- May not be able to coordinate their suck, swallow, and breathing reflexes, causing difficulties with sustaining breastfeeding or even bottle feeding.
- Are more easily fatigued or overstimulated.

Patience is needed as your baby has to learn abilities before they have completely matured. Skin to skin is very important with helping to awaken feeding behaviors, and regulating breathing, temperature and blood sugars.

Late preterm babies will need to be awakened every 3 hours for feedings. They are more likely to require supplemental feedings of pumped breast milk or formula. Pumping after breastfeeding often is recommended to help establish a plentiful milk supply.

MILK COLLECTION AND STORAGE

When mom and baby are separated due to baby's admission to NICU or SCN, pumping should be initiated within 6 hours of birth.

PUMPING GUIDELINES:

- Use a medical/hospital grade breast pump to initiate your milk supply.
- Wash your hands before pumping.
- Pump both breasts (double pumping) at the same time:
- Double pumping increases the amount of the hormone prolactin and will help to stimulate and increase your milk supply

PUMP SETTINGS:

- Pump on maximum comfortable suction – if the suction is too strong or painful turn it down.
- Pump cycles - set cycles to maximum to stimulate supply. Once milk is in - adjust to comfort, often slower cycles and more suction helps milk let down better.
- Pumping should not be uncomfortable even on low setting. If pumping is uncomfortable let your Lactation Consultant or nurse know. You may need a different flange size.
- Pump 8-10 times in 24 hours (about every 2 to 3 hours) *
- Pump a minimum of 1 time between midnight/12 am and 5 am.
- Pump for 15 minutes each time you pump, using the collection containers provided.
- All pumped breast milk must be labeled with name, date, and time milk collected.

(These instructions may be changed by your lactation consultant based upon your medical condition*.)

“It is NORMAL not to see any expressed colostrum/breast milk for the first 2 to 3 days” Keep Pumping!

After each use, rinse your breast pump kit pieces that touch your breast or come in contact with milk in cold water (flanges, valve/filter and bottle). Then wash with hot soapy water in a basin. Rinse with hot water and dry with a paper towel or place kit pieces to air dry on a towel. Do not wash or get the tubing wet. You can use a dishwasher per the manufacturer guidelines. For more specific instruction, go to www.cdc.gov and refer to instructions on “how to keep your breast pump kit clean.”

It is recommended to sanitize your pump parts at least once daily per the CDC. After the pump kit has been cleaned it can be sanitized using steam, boiling water or in a dishwasher with a sanitizing setting. It is particularly important to sanitize pump pieces if your baby is less than 3 months old, was born prematurely, or has a weakened immune system due to illness or medical treatment. For more specific instructions on sanitizing go to www.cdc.gov/healthywater/hygiene/healthychildcare/infantfeeding.html.

RENTING A BREAST PUMP:

Refer to your local Lactation Consultant for a list a breast pump rental stations. Check with your insurance company regarding what breast pump coverage you may have.

Breast pumps may be available through WIC for mothers who qualify. Types of pumps: Breast pumps are not all equal in quality and use.

- Hospital grade/rental pumps are best for a hospitalized premature or ill infant or when a mom is exclusively pumping. Using a hospital grade rental pump is the best option. Studies show these pumps establish a better and more plentiful milk supply. Hospital grade pumps have a larger, stronger motor and are designed to be used around the clock every 2-3 hours.
- Personal use double electric pumps are generally used for employed mothers who will be pumping while at work; and are not designed to be used around the clock every 2-3 hours, as the motor is not strong enough.
- Single electric or battery pumps are designed for occasional, not daily use. They take twice the time to use because each breast is pumped separately.
- Manual/hand operated pumps are also best for occasional use.

USED/BORROWED BREAST PUMPS ARE NOT HYGIENICALLY SAFE AND ARE NOT RECOMMENDED

TIPS FOR INCREASING MILK PRODUCTION WHEN PUMPING:

- If able, do skin to skin or kangaroo care in nursery and pump at baby's bedside, helps stimulate hormones.
- Breast massage and hand expression can also help stimulate hormones and help with let down.
- Make sure you are drinking to your thirst, and eating nutritious foods.

SIMPLE/EASY BREAST MILK STORAGE GUIDELINES

In the hospital setting, please make sure your freshly pumped breast milk is stored in the breast milk refrigerator in the Nursery within four hours of pumping. Only milk properly labeled with name and date and time pumped can be accepted.



(If your baby was born prematurely or is hospitalized, check with your healthcare provider for guidelines for your specific situation. Refer to www.cdc.gov for further breastmilk storage guidelines.)

Human Milk Storage Guidelines			
STORAGE LOCATIONS AND TEMPERATURES			
TYPE OF BREAST MILK	Countertop 77°F (25°C) or colder <i>(room temperature)</i>	Refrigerator 40 °F (4°C)	Freezer 0°F (-18°C) or colder
Freshly Expressed or Pumped	Up to 4 Hours	Up to 4 Days	Within 6 months is best Up to 12 months is acceptable
Thawed, Previously Frozen	1–2 Hours	Up to 1 Day <i>(24 hours)</i>	NEVER refreeze human milk after it has been thawed
Leftover from a Feeding <i>(baby did not finish the bottle)</i>	Use within 2 hours after the baby is finished feeding		

These guidelines are for healthy full-term babies and may vary for premature or sick babies. Check with your health care provider.

Find more breastfeeding resources at: WICBreastfeeding.fns.usda.gov
www.cdc.gov/breastfeeding/

MEDICALLY INDICATED REASONS TO SUPPLEMENT:

- Infant's low blood sugar.
- Inadequate urine output/dehydration.
- Weight loss of 10% of birth weight.
- Jaundice.
- Infant with temperature instability.
- Late Pre-term Infant.

If your infant requires supplementation, ask your nurse or lactation consultant about alternatives to bottle feeding: Spoon or cup feeding.

- Finger feeding with a periodontal syringe.
- Supplemental feeding system (SNS).
- Paced bottle feeding.

Try hand/manual expression of your breasts (see pages 22-23) or ask to use a breast pump so baby can be supplemented with your colostrum or breast milk.

GUIDELINES FOR COMPLEMENTARY FEEDINGS (OFFERING SOMETHING AFTER BABY HAS BEEN AT BREAST)

- Day of life 1: 5 - 7 mL
- Day of life 2: 10 -15 mL
- Day of life 3: 22 -27 mL

Supplementary feedings are designed to replace breastfeeding for those mothers who are unable to breastfeed. The volume is based on a newborn's stomach capacity.

Complementary feedings should be given by an alternative feeding method. Please see pages 42-43.

BLOOD SUGAR MONITORING

There are several circumstances that can result in your baby's pediatrician to order blood sugar monitoring for your baby. Low blood sugars can be dangerous for newborns, even resulting in seizures if gets too low. Blood sugars are tested by collecting a small drop of blood with a heel prick. Some of the reasons for this monitoring are:

- Infants of diabetic mothers. Because of the higher insulin levels a baby produces when the mother has diabetes, once born, these babies can have unstable blood sugars.
- Large for gestational age babies (typically over 9 pounds) because these babies may burn and also need more calories to stay warm.
- Small for gestational age/low birth weight babies (typically less than 6 lbs) because they have less fat stores, also may burn/need more calories to stay warm.
- Late pre-term babies (born between 35 and 37 weeks):
 - Often are smaller, less fat stores.
 - More sleepy, less energy and stamina to breastfeed well.
 - Sometimes have poor sucking skills due to immaturity

All of the above babies may need blood sugar monitoring for 12-24 hours or as ordered by your pediatrician. For low blood sugars, supplementation with expressed breast milk or formula may be required for your infant's health and safety.

Your nurse should let you know what is considered low blood sugar and help you with your infant's feeding plan as needed. Blood sugars that do not respond to your feeding plan may result in your infant being admitted to the SCN/NICU for IV glucose.

Some tips for helping decrease the risk for low blood sugars:

- Keep baby warm and skin on skin.
- Delay giving a bath until it is known sugars are stable as bathing can stress baby and lower temperature, therefore lowering blood sugars.
- Use awakening techniques for sleepy babies.
- Hand express colostrum into baby's mouth if baby's not interested in breastfeeding.

Reduce risk of nipple preference by supplementing using one of the following alternative feeding methods.

ALTERNATIVE METHODS OF SUPPLEMENTATION

PACE BOTTLE FEEDING

Pace bottle feeding is a supplemental method of feeding that can help reduce the risk of nipple confusion in baby's that are breast/bottle feeding.

1. Wash hands, gather equipment - bottle with an appropriate nipple (preferably slow flow nipple) and expressed breast milk or formula
2. Place swaddled infant in an upright position and hold bottle at a 90 degree angle.
3. Introduce nipple to baby and slowly place into mouth.
4. Every 2-3 sucks tilt bottle downward at a 45 degree angle to give baby a pause in the flow of milk. Care must be taken to prevent the baby from swallowing too much air.

5. Remove the nipple or slip it out while the tip touches infants lower lip every 6-10 sucks in order to give a rest of 3-5 seconds between sucking bursts. This method should not be used with baby's with a disorganized suck as it can make it worse.

SPOON FEEDING

1. Wash hands and plastic spoon in warm soapy water and rinse well.
2. Hand express (see section in the Guide for Successful Breastfeeding booklet on hand expression) colostrum with plastic spoon under nipple to collect droplets of colostrum.
3. Place spoon at baby's lips so that baby will slurp, sip or lick colostrum from spoon.
4. Alternatively, mom can place colostrum on her finger and have baby suck from finger. Wash the spoon in warm soapy water after each use.
5. Record the time and how many spoonfuls your infant took.

CUP FEEDING

Cup feeding best simulates the tongue and jaw movements like when baby breastfeeds.

1. Wash hands and gather equipment - expressed breast milk or formula and cup (medicine cup may be used).
2. Fill cup with expressed breast milk or formula.
3. Swaddle baby in blanket and hold baby in upright position.
4. DO NOT POUR MILK INTO BABY'S MOUTH.
5. Put cup to baby's lower lip, tip cup slightly so that milk touches upper lip. Allow baby to slurp, sip or lap liquid with tongue. Allow baby to pace feed allowing time to swallow. Leave cup at lip for entire feed.
6. Refill cup as needed, once cup becomes 1/2 empty to avoid having to tilt too far.
7. Burp baby from time to time.
8. Clean the cup in warm soapy water after each use.
9. Record amount and the time infant ate.

SYRINGE FEEDS (FINGER FEEDING)

Syringe/finger feeds are used to feed baby's with a poor dysfunctional and/ or uncoordinated suck as part of suck training.

1. Gather equipment - periodontal/curve tip syringe, expressed breast milk or formula.
2. Wash your hands. Make sure your fingernails are short and clean.
3. Fill syringe with expressed breast milk or formula.
4. Swaddle baby, place in semi-upright position in arms or lap.
5. Slide finger into baby's mouth about an inch to an inch and a half with nail bed resting on baby's tongue. Gently massage the roof of baby's mouth about half an inch.
6. After baby sucks 3-4 times, slide the curved section of the syringe tip along the side of finger into corner of baby's mouth about half an inch.
7. Gently compress plunger slowly when baby sucks. **WHEN BABY STOPS SUCKING STOP PRESSING PLUNGER.**
8. Do not press plunger faster than baby sucks and swallows. **DO NOT PRESS THE PLUNGER HARD IF PLUNGER STICKS.** It may force too much milk, and baby may choke.
9. Tickle roof of baby's mouth with your finger, not the syringe if baby stops sucking.
10. If baby holds tongue behind his lower gums, place slight downward pressure on back of tongue. This is necessary for proper breast feeding.
11. Burp baby after each syringe of milk. Record the time baby was fed and amounts.
12. Clean syringe - take syringe apart washing in hot water with mild liquid soap. Rinse well and air dry.



WEANING

When to start weaning is a personal decision. Some infants may start the process around six months when solid foods are introduced. Some mothers set a goal to wean by one year. However, many babies love to breastfeed and may continue morning, nap time, and bedtime feeds until two or older. Whatever the reason, the key to weaning success is to make it a slow gradual process for both mom and baby.

- If your infant is nursing (or you are pumping) eight times per day, take away one feeding (or pump session) and hold off nursing or pumping until next feeding.
- Use ice packs for any discomfort or relief of swelling.
- Continue the process gradually, taking away another feeding or pump session every 2-3 days (or your body is comfortable withholding the breastfeeding or pump session).
- As your body gets the message of decreased emptying, milk production slows.
- If at any time, you get uncomfortable holding milk and tenderness is not relieved by ice packs, hand express or pump just enough milk to make yourself comfortable. Be careful not to empty breast completely, or this will encourage more milk production.
- When ready to completely stop breastfeeding, you may use cabbage leaves to help the drying process. Apply clean refrigerated leaves of green cabbage next to the breast tissue under bra and change when wilted. May apply ice packs over cabbage for added relief.
- Watch for painful lumps/clogged milk ducts during the weaning process. Massage tender lumps while nursing or pumping to help express milk. Apply cabbage/ice packs treatment for relief.
- Some women find non-steroidal anti-inflammatory medicines, such as ibuprofen, to be helpful during the weaning process to control inflammation. Check with your care provider as needed to determine if this is a good option for you.
- Sunflower Seed Lecithin is a vitamin supplement found in vitamin and health food stores or on-line. It is used as an emulsifier, added in foods to help ingredients to mix more easily. In breastfeeding, it can help during engorgement, plugged ducts, mastitis, and weaning to prevent and ease those problems.
- Contact your Lactation Consultant or care provider if you develop a fever above 101 Fahrenheit. Or flu-like symptoms, or plugged ducts or discomfort that is not relieved by the suggestions listed above.

ADDITIONAL RESOURCES

WIC (WOMEN, INFANTS AND CHILDREN)

If you qualify for WIC, you have access to all of the breastfeeding services they provide. WIC has breastfeeding peer counselors, electric pumps available for mothers of infants still hospitalized, classes and various support groups. Call your WIC office for further program details, or call your local Health Department to sign up or to see if you qualify.

- Mom’s Doctor and phone number
- Baby’s Doctor and phone number
- Baby’s first appointment*
- Baby’s birth weight
- Baby’s discharge weight

*Sometimes the first “well baby exam” is scheduled for two weeks after birth, however, it is recommended that your baby has at least a weight check done during the first week.

INTERNET RESOURCES

- www.ilca.org
- www.laleche.org
- [womenshealth.gov](http://www.womenshealth.gov)
- cdc.gov
- www.lowmilksupply.org
- www.bfar.org(breastfeeding after reduction surgery)
- [www.lactationtraining.com/resources/parents handouts](http://www.lactationtraining.com/resources/parents%20handouts)
- www.kellymom.com
- toxnet.nlm.nih.gov
- aap.org

REFERENCES

1. Childbirth Graphics, Ltd. (1989). *Pocket Guide for Successful Breastfeeding*. Illustrations by Jean K. Stephens. Childbirth Graphics, Ltd.
2. Huggins, Kathleen (2005). *The Nursing Mother's Companion, (5th ed.)*. The Harvard Common Press.
3. Lawrence, Ruth A. (2005). *Breastfeeding: A Guide for the Medical Profession, (6th ed.)*. The C.Up. Mosby Company.
4. LaLeche League International (2004). *The Womanly Art of Breastfeeding (7th rev. ed.)*. A Plume Book New American Library.
5. Hale and Rowe, Hilary E., Thomas W. (2014). *Medications and Mothers' Milk (14th ed.)*. Hale Publishing LP.
6. Riordan, Jan & Auerbach, Kathleen (2005). *Breastfeeding and Human Lactation, (3rd ed.)*, Jones & Bartlett.
7. 1nJoy Birth and Parenting Videos, Inc. (2008). *Better Breastfeeding*. See What You Read series.
8. Newman, Jack (2009). *The Importance of Skin to Skin Contact*. CanadianBreastfeedingFoundation.org.
9. Mohrbaeher, Nancy and Kendal-Tackett, Kathleen (2005). *Breastfeeding Made Simple: The Seven Natural Laws of Breastfeeding*. New Harbinger Publications Inc.
10. Smilie, Christine M., MD (2007). *Baby Led Breastfeeding: The Mother-Baby Dance*. Geddes Productions.
11. Hertz, Gail S., MD/BCLC (2007). *The Little Green Book of Breastfeeding Management (4th ed.)*. Pocket Publications.
12. Ameda Belly Ball Lactation Education Tool adapted from Linda J. Smith's (2002) *Coach's Notebook: Games and Strategies for Lactation Education*. Jones and Bartlett.
13. Mohrbacher, Nancy (2010). *Breastfeeding Answers Made Simple: A Guide for Helping Mothers*. Hale Publishing LP.
14. Hughes, Vergie. www.lactationtraining.com
15. Spangler, Amy (2012). *Breastfeeding: Keep It Simple*. Baby GooRoo Publishing.
16. <http://www.bfmed.org/Resources/Protocols.aspx>

NAME: _____

PHONE: _____

Please call the hospital breastfeeding resource and/or your OB or pediatrician for any breastfeeding problems you may encounter, including:

- 3–5 day old sleepy baby that will not eat;
- Baby wetting less than 6 diapers and/or no stools in 24 hour period (*only a concern after the baby is 5 days old*);
- Baby accepts bottle but not breast;
- Mother experiencing cracked nipples beyond 2 weeks postpartum;
- Severe engorgement;
- Baby is fussy and you are concerned.
- Signs of mastitis: fever above 101, flu-like symptoms, painful breasts, breasts that are hard and hot to the touch.
- No changes in your breasts by day 5

Call 911

Baby Symptoms:

Difficulty waking baby
Difficulty breathing, choking
Bright red bleeding from circumcision that will not stop

Call Your Doctor

Baby Symptoms:

Temperature below 97 F or above 100.4 F
No wet diapers within 24 hours
Persistent vomiting
Refuses feedings
Bright red bleeding from umbilical stump
Foul odor or drainage from umbilical stump
Any rash that does not clear up
Signs of jaundice (yellow eyes/skin)
If you believe your baby to be in pain

BABY'S SECOND NIGHT

You've made it through your first 24 hours as a new mom. Maybe you have other children, but you are a new mom all over again....and now its your baby's second night.

All of a sudden, your little one discovers that he's no longer back in the warmth and comfort – albeit a bit crowded – womb where he has spent the last 8 ½ or 9 months – and it is SCARY out here! He isn't hearing your familiar heartbeat, the swooshing of the placental arteries, the soothing sound of your lungs or the comforting gurgling of your intestines. Instead, he's in a crib, swaddled in a diaper, a tee-shirt, a hat and a blanket. All sorts of people have been handling him, and he's not yet become accustomed to the new noises, lights, sounds and smells. He has found one thing though, and that's his voice....and you find that each time you take him off the breast where he comfortably drifted off to sleep, and put him in the bassinet – he protests, loudly!

In fact, each time you put him back on the breast he nurses for a little bit and then goes to sleep. As you take him off and put him back to bed – he cries again....and starts rooting around, looking for you. This goes on – seemingly for hours. A lot of moms are convinced it is because their milk isn't "in" yet, and the baby is starving. However, it isn't that, but the baby's sudden awakening to the fact that the most comforting and comfortable place for him to be is at the breast. It's the closest to "home" he can get. It seems that this is pretty universal among babies – lactation consultants all over the world have noticed the same thing.

So, what do you do? When he drifts off to sleep at the breast after a good feed, break the suction and slide your nipple gently out of his mouth. Don't move him except to pillow his head more comfortably on your breast. Don't try and burp him – just snuggle with him until he falls into a deep sleep where he won't be disturbed by being moved. Babies go into a light sleep state (REM) first, and then cycle in and out of REM and deep sleep about every ½ hour or so. If he starts to root and act as though he wants to go back to breast, that's fine...this is his way of settling and comforting. During deep sleep, the baby's breathing is very quiet and regular, and there is no movement beneath his eyelids.

Another helpful hint....his hands were his best friends in utero...he could suck on his thumb or his fingers anytime he was the slightest bit disturbed or

uncomfortable. And all of a sudden he's had them taken away from him and someone has put mittens on him! He has no way of soothing himself with those mittens on. Babies need to touch – to feel – and even his touch on your breast will increase your oxytocin levels which will help boost your milk supply! So take the mittens off and loosen his blanket so he can get to his hands. He might scratch himself, but it will heal very rapidly – after all, he had fingernails when he was inside you, and no one put mittens on him then!

FIRST TWO WEEKS: DAILY BREASTFEEDING LOG

- Fill in the hour above the box each time your baby nurses.
- Circle the W when your baby has a wet diaper.
- Fill in the number of minutes on each breast inside the box.

GOAL: 8 TO 12 FEEDINGS EVERY 24 HOURS.

- Day 1 remember it is expected that newborns do not have a good appetite and most would rather sleep and snuggle than eat. Wake baby every 3 hours to offer breast, but don't force. Your baby's appetite will perk up after the first 24 hours.

Day 1	HOUR:								
Day 2	HOUR:								
Day 3	HOUR:								
Day 4	HOUR:								
Day 5	HOUR:								

By the way – this might happen every once in a while at home too, particularly if you’ve changed his environment such as going to the doctor’s, to church, to the mall, or to the grandparents! Don’t let it throw you – sometimes babies just need some extra snuggling at the breast, because for the baby, the breast is “home.”

© 2012/Lactation Education Consultants
 May be copied and distributed freely. May not be sold.

- Circle the S when your baby has a soiled diaper.
- The days start from time of birth so if your baby is born at 8pm, day 2 won’t begin until 8pm the next day.
- It’s okay if your baby has more wet or soiled diapers than indicated here. You should call your doctor if your baby has less than the number on this log. If you cannot tell that the diaper is wet, do not count it. Your baby should have at least 1-2 stools, with all stools totaling at least a palm size amount, in a 24-hour period. See page 24 for signs that the baby is getting enough to eat.

/	/	/	/	WET DIAPERS BLACK OR BROWN TARRY SOILED DIAPERS:	W S				
/	/	/	/	WET DIAPERS BLACK OR BROWN TARRY SOILED DIAPERS:	W S	W S			
/	/	/	/	WET DIAPERS BROWNISH-GREEN TO YELLOW SOILED DIAPERS:	W S	W S	W		
/	/	/	/	WET DIAPERS GREEN/YELLOW SOILED DIAPERS:	W S	W S	W	W	W

Day 6	HOUR:								
Day 7	HOUR:								
Day 8	HOUR:								
Day 9	HOUR:								
Day 10	HOUR:								
Day 11	HOUR:								
Day 12	HOUR:								
Day 13	HOUR:								
Day 14	HOUR:								

				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W
				WET DIAPERS YELLOW SOILED DIAPERS:	W S	W S	W	W	W	W

NOTES:

A series of horizontal dotted lines for writing notes.

NOTES:

A series of horizontal dotted lines for writing notes.



SENTARA®

sentara.com/maternity

Atención: si habla español, tiene a su disposición servicios lingüísticos gratuitos. Llame al 844-809-6648.

주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 844-809-6648 번으로 전화해 주십시오.

注意: 如果您讲中文普通话, 则将为您提供免费的语言辅助服务。请致电 844_809_6648。

***ATTENTION: Language assistance services are available to you free of charge.
Call 844-809-6648***

Sentara Healthcare complies with applicable Federal Civil Rights Laws and does not exclude, deny benefits to, or otherwise discriminate against any person on the grounds of race, culture, color, religion, marital status, age, sex, sexual orientation, gender identity or gender expression, national origin or any disability or handicap.