SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; (<u>Pharmacy</u>) 1-800-750-9692. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: (Select drug below) ☐ deferasirox (Exjade®) tablets for oral	
suspension	□ deferasirox (Jadenu®) tablets
□ deferasirox (Jadenu® sprinkle) packet	
MEMBER & PRESCRIBER INFORMATI	ON: Authorization may be delayed if incomplete.
Member Name:	
Member Sentara #:	
Prescriber Name:	
Prescriber Signature:	
Office Contact Name:	
Phone Number:	
NPI #:	
DRUG INFORMATION: Authorization may be	
Drug Form/Strength:	
Dosing Schedule:	
Diagnosis:	ICD Code, if applicable:
Weight (if applicable):	Date weight obtained:
Recommended Dosing Limits:	
• deferasirox (Exjade): Maximum of 40 mg/kg/c	•
• deferasirox (Jadenu): Maximum of 28 mg/kg/	day.
CLINICAL CRITERIA: Check below all that apsupport each line checked, all documentation, including provided or request may be denied. Check the diagnost	g lab results, diagnostics, and/or chart notes, must be
☐ For diagnosis of transfusional iron overlo	oad (transfusional hemosiderosis)

(Continued on next page)

Initial Authorization: 6 months

	Member is \geq 2 years of age and has a diagnosis of transfusional hemosiderosis (i.e., transfusion of \geq 100 mL/kg of packed red blood cells, approximately 20 units for a 40 kg patient)
	Member's serum ferritin levels are consistently >1,000 mcg/L (submit serum ferritin labs done within the last 30 days)
	Medication will be dosed according to FDA recommendations for age, weight, eGFR and serum ferritin levels
	According to the prescriber, member meets BOTH of the following:
	☐ Member has an eGFR $\ge 40 \text{ mL/min}/1.73 \text{ m}^2$
	☐ Member's baseline liver function labs have been evaluated (i.e., ALT, AST, bilirubin)
	If requesting brand Jadenu or Exjade, documentation of trial and intolerable life-endangering adverse event with generic deferasirox must be submitted
□ Fo	or diagnosis of non-transfusion-dependent thalassemia syndrome
<u>Initia</u>	al Authorization: 6 months
	Member is ≥10 years of age and has a diagnosis of non-transfusion-dependent thalassemia syndrome
	Liver iron concentration (LIC) is ≥ 5 mg of Fe/g of dry weight (submit current liver biopsy, MRI or other FDA-approved test to document LIC)
	Serum ferritin is $> 300 \text{ mcg/L}$ (submit 2 serum ferritin labs, taken at least 1 month apart, from within the last 3 months)
	Medication will be dosed according to FDA recommendations for age, weight, eGFR, LIC and serum ferritin levels
	According to the prescriber, member meets <u>ALL</u> the following:
	□ Member has an eGFR ≥ 40 mL/min/1.73 m2
	☐ Member's platelets levels are $\geq 50 \times 10^9/L$
	☐ Member's baseline liver function labs have been evaluated (i.e., ALT, AST, bilirubin)
	If requesting brand Jadenu or Exjade, documentation of trial and intolerable life-endangering adverse event with generic deferasirox must be submitted
be me	thorization: 12 months. ALL DIAGNOSES. Check below all that apply. All criteria must et for approval. To support each line checked, all documentation, including lab results, diagnostics, r chart notes, must be provided or request may be denied.
	Serum ferritin has decreased from baseline or last approval (submit current ferritin labs)
	If serum ferritin is < 500 mcg/L or LIC is less than 3 mg Fe/g dw, deferasirox therapy will be temporarily discontinued; if < 300 mcg/L, deferasirox therapy will be interrupted and LIC obtained

(Continued on next page)

PA deferaxirox (Exjade)_(Jadenu) (CORE) (continued from previous page)

	According to the prescriber, member meets <u>ALL</u> the following:		
	☐ Member's liver function continues to be monitored (i.e. ALT, AST, bilirubin labs)		
	☐ Member's platelets levels are $\ge 50 \times 10^9$ /L		
	☐ For diagnosis of non-transfusion-dependent thalassemia syndrome: Member's liver iron concentration (LIC) continues to be monitored according to guideline recommendations		
	If requesting brand Jadenu or Exjade, documentation of trial and intolerable life-endangering adverse event with generic deferasirox must be submitted		
1edi	Iedication being provided by Specialty Pharmacy – Proprium Rx		

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *