SENTARA COMMUNITY PLAN
(MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: **The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request**. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Skyrizi™ (risankizumab-rzaa) **Injection**

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ **Date of Birth:** _____

Prescriber Name: _____

Prescriber Signature: _____ **Date:** _____

Office Contact Name: _____

Phone Number: _____ **Fax Number:** _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ **Length of Therapy**: _____

Diagnosis: _____ **ICD Code, if applicable:** _____

Weight (if applicable): _____ **Date weight obtained:** _____

Requested Medication is ONE of the following:

- ❑ New Start
- ❑ Continuation of therapy – Provide therapy start date:_____

DIAGNOSIS	RECOMMENDED DOSE/ QUANTITY LIMIT
Plaque Psoriasis/Psoriatic Arthritis	• Dosage 150mg Pen or Syringe (one injection) administered by subcutaneous injection at Week 0, Week 4 and every 12 weeks thereafter • Quantity Limit: • Two, 150 mg syringes or pen allowed in the initial 28 days • One, 150mg pen/ syringe per 84 days after induction period • Pediatrics: • Weight < 40kg: 55mg pen/syringe (one injection) administered by subcutaneous injection at week 0, week 4, and every 12 weeks thereafter • Weight ≥ 40 kg: 150 mg pen/syringe (one injection) administered by subcutaneous injection at week 0, week 4 and every 12 weeks thereafter

(Continued on next page)

DIAGNOSIS	RECOMMENDED DOSE/ QUANTITY LIMIT
Crohn's Disease/Ulcerative Colitis (UC)	• IV loading dose 600mg (1200mg for UC) at weeks 0, 4 and 8. Then via subq prefilled cartridge: 180 to 360 mg at week 12 and every 8 weeks thereafter; use lowest effective dosage to maintain therapeutic response • Quantity Limit: • One, 180 or 360mg pen/ syringe per 56 days after induction period

ATTENTION: Skyrizi IV loading dose for treatment of Crohn's disease and Ulcerative colitis can only be billed under the **MEDICAL BENEFIT**. NDC: 00074-5015-01; J2327; 600mg= 600 billable units, 1200mg= 1200 billable units

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Authorization Length : 12 months

❑ DIAGNOSIS: Moderate-to-Severe Chronic Plaque Psoriasis

- ❑ Member is 6 years of age or older
- ❑ Member has a diagnosis of moderate to severe plaque psoriasis for ≥ 6 months with ≥ 1 of the following:
  - ❑ Affected body surface area (BSA) of ≥ 10%
  - ❑ Psoriasis Area and Severity Index (PASI) score ≥ 10
  - ❑ Incapacitation due to plaque location (head and neck, palms, soles or genitalia)
- ❑ Member is not receiving risankizumab-rzaa in combination with another biologic agent for psoriasis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib)
- ❑ Trial and failure of **TWO (2)** preferred drugs below:

❑ adalimumab-adbm (Boehringer Ingelheim) **OR** Hadlima® (adalimumab-bwwd)	❑ Enbrel®	❑ Pyzchiva® syringe/vial **OR** Starjemza™ (Requires trial and failure of a preferred TNF-alpha inhibitor)

❑ DIAGNOSIS: Psoriatic Arthritis

- ❑ Member has a diagnosis of active psoriatic arthritis
- ❑ Member is 6 years of age or older
- ❑ Member is not receiving risankizumab-rzaa in combination with another biologic agent for psoriatic arthritis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib)
- ❑ Trial and failure of **TWO (2)** preferred drugs below:

❑ adalimumab-adbm (Boehringer Ingelheim) **OR** Hadlima® (adalimumab-bwwd)	❑ Enbrel®	❑ Pyzchiva® syringe/vial **OR** Starjemza™ (Requires trial and failure of a preferred TNF-alpha inhibitor)

(Continued on next page)

❑ DIAGNOSIS: Crohn's Disease

- ❑ Member has a diagnosis of moderate-to severely active Crohn's Disease
- ❑ Member is 18 years of age or older
- ❑ Trial and failure of a compliant regimen of corticosteroids or immunomodulators (e.g. azathioprine, 6-mercaptopurine, or methotrexate)
- ❑ Member is not receiving risankizumab-rzaa in combination with another biologic agent for Crohn's disease or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib, upadacitinib)
- ❑ Trial and failure of **BOTH** of the preferred drugs below:

❑ adalimumab-adbm (Boehringer Ingelheim) **OR** Hadlima® (adalimumab-bwwd)	❑ Pyzchiva® syringe/vial **OR** Starjemza™ (Requires trial and failure of a preferred TNF-alpha inhibitor)

❑ DIAGNOSIS: Ulcerative Colitis

- ❑ Member has a diagnosis of moderate-to severely active Ulcerative Colitis
- ❑ Member is 18 years of age or older
- ❑ Member is not receiving risankizumab-rzaa in combination with another biologic agent for ulcerative colitis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib, upadacitinib)
- ❑ Trial and failure of **BOTH** of the preferred drugs below:

❑ adalimumab-adbm (Boehringer Ingelheim) **OR** Hadlima® (adalimumab-bwwd)	❑ Pyzchiva® syringe/vial **OR** Starjemza™ (Requires trial and failure of a preferred TNF-alpha inhibitor)

❑ Induction Dose (If required) – One time approval for duration of 2 months, member to receive up to three (3) IV infusion doses

Authorization Criteria: To be reviewed for one-time approval under the medical benefit

- ❑ Medication will be used as induction therapy
- ❑ Medication being provided by:

  Location/site of drug administration: _____

  NPI or DEA # of administering location: _____

- ❑ Member to receive FDA approved loading dose for **ONE** of the following indications:
  - ❑ Crohn's Disease- 600mg administered by IV infusion over a period of at least one hour at week 0,4 and 8
  - ❑ Ulcerative Colitis: 1200mg administered by IV infusion over a period of at least one hour at week 0, 4 and 8

(Continued on next page)

Medication being provided by Specialty Pharmacy – PropriumRx

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.*

REVISED/UPDATED/REFORMATTED: ~~12/17/2019; 11/23/2020; 7/19/2021;1/5/2022; 7/11/2022; 6/26/2023; 12/29/2023; 1/18/2024, 8/27/2024, 12/13/2024; 3/25/2025, 10/17/2025, 12/15/2025; 1/29/2026, 2/24/2026; 3/11/2026; 7/13/2026, 8/6/2026;~~ **9/30/2026**