

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

**Drug Requested:** Skyrizi™ (risankizumab-rzaa) Injection

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

Diagnosis	Recommended Dose/ Quantity Limit
<input type="checkbox"/> <b>Plaque Psoriasis/Psoriatic Arthritis</b>	<ul style="list-style-type: none"><li>• Dosage 150mg Pen or Syringe (one injection) administered by subcutaneous injection at Week 0, Week 4 and every 12 weeks thereafter</li><li>• Quantity Limit:</li><li>• Two, 150 mg syringes or pen allowed in the initial 28 days</li><li>• One, 150mg pen/ syringe per 84 days after induction period</li></ul>
<input type="checkbox"/> <b>Crohn's Disease/Ulcerative Colitis (UC)</b>	<ul style="list-style-type: none"><li>• IV loading dose 600mg (1200mg for UC) at weeks 0, 4 and 8. Then via subq prefilled cartridge: 180 to 360 mg at week 12 and every 8 weeks thereafter; use lowest effective dosage to maintain therapeutic response</li><li>• Quantity Limit:</li><li>• One, 180 or 360mg pen/ syringe per 84 days after induction period</li></ul>

(Continued on next page)

**ATTENTION:** Skyrizi IV loading dose for treatment of Crohn's disease and Ulcerative colitis can only be billed under the **MEDICAL BENEFIT**. NDC: 00074-5015-01; J2327; 600mg= 600 billable units, 1200mg= 1200 billable units

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**❑ DIAGNOSIS: Moderate-to-Severe Chronic Plaque Psoriasis.**

- ☐ Member is 18 years of age or older
- ☐ Member has a diagnosis of moderate to severe plaque psoriasis for  $\geq 6$  months with  $\geq 1$  of the following:
  - ☐ Affected body surface area (BSA) of  $\geq 10\%$
  - ☐ Psoriasis Area and Severity Index (PASI) score  $\geq 10$
  - ☐ Incapacitation due to plaque location (head and neck, palms, soles or genitalia)
- ☐ Member did not respond adequately (or is not a candidate) to a 3-month minimum trial of topical agents (anthralin, coal tar preparations, corticosteroids, emollients, immunosuppressives, keratolytics, retinoic acid derivatives, and/or Vitamin D analogues)
- ☐ Member did not respond adequately (or is not a candidate) to a 3-month minimum trial of  $\geq 1$  systemic agent (e.g., immunosuppressives, retinoic acid derivatives, and/or methotrexate)
- ☐ Member did not respond adequately (or is not a candidate) to a 3-month minimum trial of phototherapy (e.g., psoralens with UVA light (PUVA) or UVB with coal tar or dithranol)
- ☐ Member is not receiving risankizumab-rzaa in combination with another biologic agent for psoriasis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib)
- ☐ Trial and failure of **TWO (2)** preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

**❑ DIAGNOSIS: Psoriatic Arthritis**

- ☐ Member has a diagnosis of moderate-to-severe psoriatic arthritis
- ☐ Member is 18 years of age or older
- ☐ Member did not respond adequately (or is not a candidate) to a 3-month minimum trial of  $\geq 1$  systemic agent (e.g., immunosuppressives, and/or methotrexate)

(Continued on next page)

- ☐ Member is not receiving risankizumab-rzaa in combination with another biologic agent for psoriatic arthritis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib)
- ☐ Trial and failure of **TWO (2)** preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

#### ☐ **DIAGNOSIS: Crohn's Disease**

- ☐ Member has a diagnosis of Crohn's Disease
- ☐ Member is 18 years of age or older
- ☐ Trial and failure of a compliant regimen of oral corticosteroids unless contraindicated or intravenous corticosteroids
- ☐ Member is not receiving risankizumab-rzaa in combination with another biologic agent for Crohn's disease or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib, upadacitinib)
- ☐ Trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Infliximab
----------------------------------	-------------------------------------

#### ☐ **DIAGNOSIS: Ulcerative Colitis**

- ☐ Member has a diagnosis of Ulcerative Colitis
- ☐ Member is 18 years of age or older
- ☐ Trial and failure to **ONE** conventional agent (i.e., 6-mercaptopurine, azathioprine, balsalazide, corticosteroids, cyclosporine, mesalamine, sulfasalazine) for at least a 3-month duration of therapy
- ☐ Member is not receiving risankizumab-rzaa in combination with another biologic agent for ulcerative colitis or non-biologic immunomodulator (e.g., apremilast, tofacitinib, baricitinib, upadacitinib)
- ☐ Trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Infliximab
----------------------------------	-------------------------------------

(Continued on next page)

☐ **Induction Dose (If required) – One time approval for duration of 2 months, member to receive up to three (3) IV infusion doses**

**Authorization Criteria: To be reviewed for one-time approval under the medical benefit**

- ☐ Medication will be used as induction therapy
- ☐ Medication being provided by:
  - ☐ **Location/site of drug administration:** \_\_\_\_\_
  - NPI or DEA # of administering location:** \_\_\_\_\_
- ☐ Member to receive FDA approved loading dose for **ONE** of the following indications:
  - ☐ Crohn's Disease- 600mg administered by IV infusion over a period of at least one hour at week 0,4 and 8
  - ☐ Ulcerative Colitis: 1200mg administered by IV infusion over a period of at least one hour at week 0,4 and 8

**Medication being provided by Specialty Pharmacy - PropriumRx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****