

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Benlysta[®] (belimumab) Subcutaneous Injection (Pharmacy)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

Quanty Limits: For adults with SLE: Maximum of 200 mg once weekly. For adults with lupus nephritis: 400 mg once weekly for 4 doses, then 200 mg once weekly thereafter

CLINICAL CRITERIA/DIAGNOSIS: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. Check box below for the Diagnosis that applies.

Diagnosis - active systemic lupus erythematosus (SLE) in adults who are receiving standard therapy

Initial Authorization: 12 months

Must be prescribed by or in consultation with a rheumatologist

(Continued on next page)

- Member is 18 years of age or older with a diagnosis of active, autoantibody-positive SLE confirmed by one of the following (**submit lab results**):
 - anti-nuclear antibody (ANA) titer \geq 1:80
 - anti-double stranded DNA (anti-dsDNA) \geq 30 IU/mL
- Member's SLE activity has been confirmed by one of the following (**submit results**):
 - Safety of Estrogen in Lupus National Assessment – Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI) score of 6-12
 - \geq 2 British Isles Lupus Assessment Group (BILAG) B organ domain scores
- Member has tried three of the following and is established on two of the following therapies taken for the last 90 days (**please submit chart notes documenting therapy trials with insufficient disease control**):

<input type="checkbox"/> mycophenolate	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> azathioprine
<input type="checkbox"/> cyclophosphamide	<input type="checkbox"/> methotrexate	<input type="checkbox"/> cyclosporine
<input type="checkbox"/> corticosteroids	<input type="checkbox"/> Other _____	

- Member does not have any of the following limitations to therapy: severe active central nervous system lupus; current or previous diagnosis of progressive multifocal leukoencephalopathy (PML); or concurrent use with other biologics

Diagnosis - active lupus nephritis in adults who are receiving standard therapy

Initial Authorization: 12 months

- Must be prescribed by or in consultation with a nephrologist or rheumatologist
- Member is 18 years of age or older with a diagnosis of active lupus nephritis Class III, IV, or V as confirmed by renal biopsy
- Member's diagnosis of active, autoantibody-positive SLE was confirmed by one of the following (**submit lab results**):
 - anti-nuclear antibody (ANA) titer \geq 1:80
 - anti-double stranded DNA (anti-dsDNA) \geq 30 IU/mL
- Member has active renal disease and has received standard therapy for the last 90 days with corticosteroids along with one of the following (**chart notes documenting established therapy must be submitted**):
 - mycophenolate
 - cyclophosphamide
- Provider must obtain a baseline measurement of one of the following collected within the last 30 days (**labs must be submitted**):
 - urine protein:creatinine ratio (uPCR)
 - urine protein

(Continued on next page)

- Member does not have any of the following limitations to therapy: severe active central nervous system lupus; current or previous diagnosis of progressive multifocal leukoencephalopathy (PML); or concurrent use with other biologics

Reauthorization Approval: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis - systemic lupus erythematosus (SLE) in adults

- All of the initial authorization criteria continues to be met
- Member's response to therapy has been confirmed by one of the following (**submit results**):
 - Safety of Estrogen in Lupus National Assessment – Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI) score has improved by and/or maintained at a level that is ≥ 4 points below baseline score
 - No new BILAG-A organ domain score OR 2 new BILAG-B organ domain scores
- Member has absence of intolerable side effects such as serious infections, signs or symptoms of progressive multifocal leukoencephalopathy (PML), malignancy, severe hypersensitivity reactions/anaphylaxis, or serious infusion reactions

Reauthorization Approval: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis - active lupus nephritis in adults

- All of the initial authorization criteria continues to be met
- Member has had improvement from baseline and/or stabilization since last approval of one of the following (**submit current labs completed within the last 30 days**):
 - Urine protein:creatinine ratio (uPCR)
 - Urine protein
- Member has absence of intolerable side effects such as serious infections, signs or symptoms of progressive multifocal leukoencephalopathy (PML), malignancy, severe hypersensitivity reactions/anaphylaxis, or serious infusion reactions

Medication being provided by a Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****