

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: (Select drug below)

<input type="checkbox"/> Oxlumo[®] (lumasiran)	<input type="checkbox"/> Rivfloza[™] (nedosiran)
--	--

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Recommended Dosage:

Oxlumo		
Body Weight	Loading Dose	Maintenance Dose
Less than 10 kg	6 mg/kg once monthly for 3 doses	3 mg/kg once monthly, beginning 1 month after the last loading dose
10 kg to less than 20 kg	6 mg/kg once monthly for 3 doses	6 mg/kg once every 3 months (quarterly), beginning 1 month after the last loading dose
20 kg and above	3 mg/kg once monthly for 3 doses	3 mg/kg once every 3 months (quarterly), beginning 1 month after the last loading dose

(Continued on next page)

Rivfloza - Recommended dosage is shown below and is administered subcutaneously once monthly			
Age	Body Weight		
	Less than 39 kg	39 kg to less than 50 kg	50 kg and above
Age 2 to less than 12 years	3.3 mg/kg	128 mg	160 mg
Age 12 years and older	128 mg		160 mg

Quantity Limits:

- Oxlumo: N/A
- Rivfloza 80 mg vial: 2 vials per month
- Rivfloza 128 mg prefilled syringe: 1 syringe per month
- Rivfloza 160 mg prefilled syringe: 1 syringe per month

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- ☐ Must be prescribed by a geneticist, nephrologist, or urologist with expertise in the diagnosis and treatment of primary hyperoxaluria type 1 (PH1)
- ☐ Member has a definitive diagnosis of PH1 confirmed by biallelic pathogenic mutation in the alanine:glyoxalate aminotransferase (AGXT) gene as identified on molecular genetic testing (**must submit documentation**)
- ☐ Member has signs and symptoms attributed to PH1 such as recurrent calcium oxalate (CaOx) kidney stones, urolithiasis, infantile oxalosis, failure to thrive and renal failure in an infant <12 months, nephrocalcinosis associated with decreased GFR, oxalate crystals in any biological fluid or tissue, increased serum creatinine with calcium oxalate (CaOx) stones, CaOx tissue deposits, renal failure of unknown causes (**must submit test results or medical chart notes confirming symptoms**)
- ☐ Provider must submit documentation to confirm member meets **ONE** of the following (**must submit lab documentation**):
 - ☐ Member had failure to achieve normalization of urinary oxalate (UOx) excretion levels after ≥ 3 months of therapy with pyridoxine (vitamin B6) dosed at 5 mg/kg/day, alkalinizing agents (potassium citrate, sodium citrate, neutral phosphate; etc.) and hyperhydration and will continue those therapies while taking the requested medication
 - ☐ Member's genetic assessment reveals a mutation consistent with pyridoxine (vitamin B6) non-responsive PH1

(Continued on next page)

- ☐ Member has **ONE** of the following (**must submit lab documentation**):
 - ☐ Increased urinary oxalate excretion (i.e. greater than 0.7 mmol/1.73 m² per day [90 mg/1.73 m² per day])
 - ☐ Increased urinary oxalate:creatinine ratio relative to normative values for age
- ☐ Member does **NOT** have severe kidney damage (eGFR <30 mL/min/1.73 m²), is **NOT** receiving hemodialysis and has **NOT** previously received a liver or kidney transplant
- ☐ Provider has submitted lab test results documenting member's current 24-hour urinary oxalate excretion (corrected for BSA): _____
- ☐ Provider has submitted lab test results documenting member's baseline plasma oxalate levels: _____
- ☐ Member does **NOT** have secondary causes of hyperoxaluria (e.g., diet with excessive intake of oxalate, gastric bypass surgery, Irritable Bowel Disease, other intestinal disorders)
- ☐ Medication will **NOT** be used in combination with other urinary oxalate reducing agents (i.e., lumasiran, nedosiran)
- ☐ Requested dosing is in accordance with the United States Food and Drug Administration approved labeling

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member continues to meet all initial authorization criteria
- ☐ Provider has submitted current medical chart notes and laboratory values documenting a positive clinical response to therapy (i.e., improvement in nephrocalcinosis, decreased formation of renal stones) as well as a clinically significant reduction from pre-treatment baseline of urinary oxalate concentration (corrected for BSA), urinary oxalate:creatinine ratio, or plasma oxalate concentrations

Medication being provided by Specialty Pharmacy – Proprium Rx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****