

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Prevmis[®] (letermovir) tablets (**Pharmacy**)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Quantity Limit: 1 tablet per day (all strengths)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis: Cytomegalovirus, prophylaxis in hematopoietic cell transplant recipients

Initiate therapy between Day 0 and Day 28 post-HSCT (before or after engraftment) and continue through Day 100 post-HSCT. In patients at risk for late CMV infection and disease, Prevmis[®] may be continued through Day 200 post-HSCT.

Recommended Dosage:

- **Adult and Pediatric Patients 12 Years of Age and Older and Weighing at least 30 kg:** 480 mg administered orally once daily

(Continued on next page)

Recommended Dosage:

- **Pediatric Patients 6 Months to Less than 12 Years of Age or 12 Years of Age and Older and Weighing Less than 30 kg:**

Body Weight	Daily Oral Dose	Tablets	Oral Pellets*
15 kg to less than 30 kg	240 mg	One 240 mg tablet	Two 120 mg packets
7.5 kg to less than 15 kg	120 mg	Not Recommended	One 120 mg packet
6 kg to less than 7.5 kg	80 mg	Not Recommended	Four 20 mg packets

*Commercial availability of oral pellets is still pending

Length of Authorization: 200 days of therapy

- Member is 6 months of age or older and weighs at least 6 kg
- Member will be receiving Prevymis® for the prophylaxis of cytomegalovirus (CMV) disease
- Member is a CMV-seropositive recipient [R+] of an allogeneic hematopoietic stem cell transplant (HSCT)
- Medication will be initiated between day 0 and day 28, before or after engraftment
 - Enter date transplant was performed: _____
- Member is **NOT** receiving the requested medication beyond 200 days post-transplantation

Diagnosis: Cytomegalovirus, prophylaxis in kidney transplant recipients

Initiate therapy between Day 0 and Day 7 post-transplant and continue through Day 200 post-transplant.

Recommended Dosage:

- **Adult and Pediatric Patients 12 Years of Age and Older and Weighing at least 40 kg:** 480 mg administered orally once daily

Length of Authorization: 200 days of therapy

- Member is 12 years of age or older and weighs at least 40 kg
- Member will be receiving a kidney transplant
- Member will be receiving Prevymis® for the prophylaxis of cytomegalovirus (CMV) disease
- Member is at high-risk for CMV disease [documentation recording kidney donor is CMV-seropositive, and the recipient (member) is CMV-seronegative (D+/R-)]
- Medication will be initiated between day 0 and day 7, before or after engraftment
 - Enter date transplant was performed: _____
- Member is **NOT** receiving the medication beyond 200 days post-transplantation

(Continued on next page)

Medication being provided by Specialty Pharmacy – Proprium Rx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.