## SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization may be delayed.</u>

**Drug Requested:** mifepristone 300 mg (Korlym®)

ME	EMBER & PRESCRIBER INFORMATION:	Authorization may be delayed if incomplete.
Meml	ber Name:	
Member Sentara #:		
	eriber Name:	
	eriber Signature:	
	e Contact Name:	
	e Number:	
	OR NPI #:	
	UG INFORMATION: Authorization may be dela	
Drug	Form/Strength:	
	ng Schedule:	
Diagn	nosis:	ICD Code, if applicable:
Weigl	ht:	Date:
Ouan	ntity Limits: 120 tablets per 30 days	
Recor	mmended Dosage: Initiate therapy with 300 mg/day ary ymptom control, and daily dose will NOT exceed 20 mg	•
suppo	INICAL CRITERIA: Check below all that apply. Fort each line checked, all documentation, including labited or request may be denied.	
Initi	ial Authorization: 6 months	
	Member is 18 years of age or older	
	Prescribing physician is an endocrinologist	
☐ Member has a diagnosis of Endogenous Cushing's Syndrome, and satisfies ONE of the Control of		endrome, and satisfies <b>ONE</b> of the following:
	<ul> <li>Diagnosis of Type 2 Diabetes Mellitus</li> <li>Glucose intolerance noted by <u>ONE</u> of the followint tolerance test or Hemoglobin A1c test (HbA1c)</li> </ul>	ng (must submit documentation): oral glucose

(Continued on next page)

	Past medical history confirms <b>ONE</b> of the following:	
	☐ Member has undergone surgery and has not been curative	
	☐ Member is <u>NOT</u> a candidate for surgery	
	Documentation of clinical failure (unable to normalize cortisol levels for the treatment of Cushing's Syndrome) to ketoconazole tablets taken along with <u>ONE</u> of the following:  An additional steroidogenesis inhibitor such as Metopirone <sup>®</sup> (metyrapone capsules) or mitotane	
	tablets	
	☐ A pituitary-directed therapy such as cabergoline or Signifor® LAR (pasireotide)	
	Documentation of clinical failure to control glucose levels with metformin AND TWO (2) of the following treatments:	
	☐ Glucagon-Like Peptide-1 (GLP-1) Receptor Agonist (e.g., Trulicity®, Ozempic®) ☐ Insulin (e.g., Humalog®, Lantus®)	
	☐ Dipeptidyl Peptidase 4 (DPP-4) Inhibitor (e.g., Januvia®, Onglyza®)	
	Member is <u>NOT</u> also taking/ will <u>NOT</u> take strong inhibitors of CYP2A medications (e.g., simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, tacrolimus)	
	For females of reproductive potential: pregnancy has been excluded before initiation of treatment and plans for prevention are implemented during treatment and for one month after stopping	
<b>Reauthorization:</b> 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.		
	Positive disease response has been achieved, demonstrated by improved glycemic control (decreased hemoglobin A1c) (current labs must be submitted to document HbA1c)	
Medication being provided by a Specialty Pharmacy – Proprium Rx		

\*\*Use of samples to initiate therapy does not meet step edit/preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. \*