

SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-844-305-2331. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If information provided is not complete, correct, or legible, authorization can be delayed.

Drug Requested: Simponi® ARIA™ (golimumab) (J-1602) (Medical) (Non-Preferred)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

- Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

DIAGNOSIS: Check diagnosis below that applies

Diagnosis – Moderate-to severe Active Rheumatoid Arthritis

Dosage: IV: 2 mg/kg at weeks 0, 4, and then every 8 weeks thereafter (in combination with methotrexate)

- Member is 18 years of age or older

(Continued on next page)

- Member has a diagnosis of moderate to severe active rheumatoid arthritis
- Must be in combination with methotrexate

OR

- Member has a contraindication or adverse reaction to methotrexate
- Trial and failure of at least **ONE (1) other DMARD** therapy (**check each tried**):

<input type="checkbox"/> methotrexate	<input type="checkbox"/> auranofin	<input type="checkbox"/> azathioprine
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> leflunomide	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> Other: _____		

- Member has trial and failure of **BOTH** preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd)	<input type="checkbox"/> Enbrel [®]
---	--

Diagnosis – Active Psoriatic Arthritis

Dosage: IV: Children ≥2 years and Adolescents: Simponi Aria: IV: 80 mg/m²/dose at weeks 0, 4, and then every 8 weeks thereafter. **Adults:** IV: 2 mg/kg at weeks 0, 4, and then every 8 weeks

- Member is 2 years of age or older
- Member has a diagnosis of psoriatic arthritis
- Member has trial and failure of **TWO (2)** preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd)	<input type="checkbox"/> Enbrel [®]	<input type="checkbox"/> Pyzchiva [®] OR Starjemza [™] syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor)
--	--	--

Diagnosis – Active Ankylosing Spondylitis

Dosage: IV: 2 mg/kg at weeks 0, 4, and then every 8 weeks thereafter

- Member is 18 years of age or older
- Member has a diagnosis of active ankylosing spondylitis
- Trial and failure of **BOTH** preferred drugs below

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima [®] (adalimumab-bwwd)	<input type="checkbox"/> Enbrel [®]
--	--

(Continued on next page)

Diagnosis – Active Polyarticular Juvenile Idiopathic Arthritis
Dosage: IV: 80 mg/m2/dose at weeks 0, 4, and then every 8 weeks thereafter

- Member is 2 years of age and older
- Member has a diagnosis of moderate to severely active juvenile rheumatoid arthritis OR juvenile idiopathic arthritis
- Member has at least five swollen joints
- Member has three or more joints with limitation of motion and pain, tenderness or both
- Member tried and failed **at least one (1)** previous **DMARD** therapy including but not limited to the following; check each tried

<input type="checkbox"/> methotrexate	<input type="checkbox"/> auranofin	<input type="checkbox"/> azathioprine
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> leflunomide	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> Other: _____		

- Trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Enbrel®
--	----------------------------------

Medication being provided by: Please check applicable box below.

- Location/site of drug administration: _____
 NPI or DEA # of administering location: _____
- OR**
- Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health’s definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member’s ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
*****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.*****