



## BEHAVIORAL HEALTH GUIDELINE

### BIPOLAR AND SCHIZOPHRENIA GUIDELINE

#### Guideline History

Date Approved	7/20
Date Revised	
Date Reviewed	3/24
Next Review Date	6/26

These Guidelines are promulgated by Sentara Healthcare (SHC) as recommendations for the clinical Management of specific conditions. Clinical data in a particular case may necessitate or permit deviation from these Guidelines. The SHC Guidelines are institutionally endorsed recommendations and are not intended as a substitute for clinical judgment.

# Medications for the Treatment of Bipolar Disorder and Schizophrenia

## MOOD STABILIZERS

Medication	Indication	Dosage	Side Effects
<b>Divalproex (Depakote)</b>			
	Bipolar I disorder- Manic	<b>DR</b> Initial 1750 mg orally daily in divided doses; increase as quickly as possible to clinical effect; usual trough plasma level 50 to 125 mcg/ml (347 to 693 mcmol/L); MAX 60 mg/kg/day  <b>ER</b> Initial 125 mg/kg/day orally once daily increase as quickly as possible to clinical effect; usual trough plasma level 85 to 125 mcg/ml (589 to 867 mcmol/L); MAX 60 mg/kg/day	Common: Alopecia, Weight gain, Abdominal pain, Weakness, Diarrhea, Nausea/Vomiting, Tremor, Headache  Serious: Hepatotoxicity, Teratogenicity, Pancreatitis, Thrombocytopenia (dose dependent)
<b>Lamotrigine (Lamictal)</b>			
	Bipolar I disorder, <i>monotherapy</i>	Initial 25mg/day orally for 2 weeks  Titration 50mg/day for 2 weeks 100mg/day for 1 week  Maintenance 200mg/day	Common: Rash, Abdominal pain, Diarrhea, Nausea/Vomiting, Dizziness, Headache, Somnolence, Blurred vision  Serious: Stevens-Johnson syndrome/Toxic Epidermal Necrolysis, Liver failure, Aseptic meningitis, Neuroleptic malignant syndrome
	Bipolar I disorder, <i>in combination with valproic acid</i>	Initial 25mg/day orally every other day for 2 weeks  Titration 25mg/day for 2 weeks 50mg/day for 1 week  Maintenance 100mg/day	
	Bipolar I disorder, <i>in combination with carbamazepine or other enzyme inducing antiepileptic</i>	Initial 50mg/day orally for 2 weeks  Titration 100mg/day for 2 weeks in divided doses 200mg/day for 1 week in divided doses 300mg/day for 1 week in divided doses  Maintenance	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

		May increase up to the usual maintenance dose of 400mg/day in divided doses	
<b>Carbamazepine ER (Equetro)</b>			
	Bipolar I disorder, acute manic or mixed episodes	<p>Initial 200mg orally twice daily</p> <p>Titration May increase in increments of 200mg/day</p> <p>Max dose Doses in excess of 1600mg/day have not been studied</p>	<p>Common: Dizziness, Drowsiness, Ataxia, Nausea, Vomiting, Pruritus, Skin Rash, Constipation, Xerostomia</p> <p>Serious: Stevens-Johnson Syndrome/Toxic Epidermal Necrolysis, Aplastic Anemia, Agranulocytosis</p>
<b>Lithium</b>			
	Bipolar I disorder, acute treatment of manic or mixed episodes <i>monotherapy</i>	<p>Immediate Release</p> <p>Initial 300mg orally 3 times daily</p> <p>Titration Add 300mg every 3 days to acute goal</p> <p>Extended Release 1800mg/day orally in 2-3 divided doses</p> <p>Target Serum Level 0.8-1.2meq/L</p>	<p>Common: Hyperparathyroidism, Weight gain, Dizziness/Fatigue, Leukocytosis, Increased thirst</p> <p>Serious: Bradycardia, Hypothyroidism, Erythema multiforme, Blurred vision, Nephrogenic diabetes insipidus</p>
	Bipolar I disorder, maintenance <i>monotherapy</i>	<p>Immediate-release tablet and capsule Initial 300mg orally 3 times daily; titrate by 300 mg every 3 days to maintenance goal, desired serum lithium level 0.8 to 1 mEq/L usually 300 to 600 mg 2 to 3 times daily.</p> <p>Extended-release tablet 900 to 1200 mg/day orally in 2 to 3 divided doses; desired serum lithium level 0.6 to 1.2 mEq/L</p>	

# Medications for the Treatment of Bipolar Disorder and Schizophrenia

## FIRST GENERATION ANTIPSYCHOTICS

Medication	Indication	Dosage	Side Effects
<b>Chlorpromazine (Thorazine)</b>			
	Bipolar Disorder, Manic Episode (hospitalized)	<p>Oral preparation:</p> <p>Initial 25mg orally 3 times a day</p> <p>Titration Increase gradually until effective dose is reached (usually 400mg/day)</p> <p>Injectable preparation:</p> <p>Initial 25mg IM for control of severe symptoms May give 25-50mg in 1 hour as needed</p> <p>Titration Increase subsequent IM doses gradually over several days as needed 400mg IM every 4-6 hours until patient is controlled</p>	<p>Common:</p> <p>Orthostatic hypotension, Tachycardia, Ineffective thermoregulation (heat stroke or hypothermia), Constipation, Xerostomia, Akathisia, Dizziness, Parkinsonism, Somnolence, Tardive dyskinesia</p> <p>Serious</p> <p>QT prolongation, Agranulocytosis, Aplastic Anemia, Anaphylaxis, Dystonia, Neuroleptic Malignant Syndrome</p>
	Schizophrenia (hospitalized)	Same as above	
	Bipolar Disorder, Maintenance	<p>Initial 10mg orally 3-4 times daily OR 25mg orally 2-3 times daily</p> <p>Titration Increase gradually until symptom control Continue optimal dosage for 2 weeks Gradually reduce to lowest effective dose</p> <p>Typical dose 200mg-800mg/day is typical target range</p>	
	Schizophrenia, Maintenance	Same as above	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

Thioridazine (Mellaril)			
	Schizophrenia, refractory	Initial 50-100mg orally 3 times a day  Titration increase gradually to a max of 800mg/day in 2-4 divided doses  Maintenance reduce dose to determine minimum maintenance dose (range 200- 800mg/day)  Max dose 800mg/day	Common: Orthostatic hypotension, Diminished sweating, Photosensitivity, Constipation, Xerostomia, Extrapyramidal disease, Somnolence, Blurred vision, Epithelial Keratopathy, Urinary Retention  Serious: QT prolongation, Paralytic Ileus, Myelosuppression, Drug induced lupus, Ineffective thermoregulation
Loxapine (Adasuve)			
	Bipolar I Disorder – psychomotor agitation	Initial/ Maintenance 10mg via oral inhalation as single dose within a 24 hour period	Common: Altered taste sense, Sedation, Pharyngitis  Serious: Bronchospasm, Cerebrovascular Accident/Transient Ischemic Attack, Death
	Schizophrenia – psychomotor agitation	Initial/ Maintenance 10mg via oral inhalation as single dose within a 24 hour period	
Perphenazine (Trilafon)			
	Schizophrenia (hospitalized)	Initial/ Maintenance 8-16mg orally 2-4 times daily  Max dose 64mg/day	Common Orthostatic hypotension, Diminished sweating, Constipation, Xerostomia, Extrapyramidal disease, Somnolence, Tardive Dyskinesia, Blurred Vision, Urinary Retention, Epithelial Keratopathy, Retinitis Pigmentosa, Nasal congestion  Serious: QT prolongation, Drug induced Lupus

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b>Fluphenazine Hydrochloride (Prolixin)</b>			
	Schizophrenia	<p>Oral preparation:</p> <p>Initial 2.5-10mg/day in divided doses every 6-8 hours and may increase to clinical effect and tolerance usually achieved at less than 20mg/day; 40mg/day may be necessary in severe cases, however safety of prolonged use is not established.</p> <p>Maintenance 1 to 5mg orally once daily after acute symptom control that is generally achieved with higher doses. Oral dose is generally 2 to 3 times the parenteral dose.</p> <p>Injection: (Acute agitation) 2 - 5mg IM Depending on the severity of symptoms, daily dosage may range from 2.5 to 10 mg IM given in divided doses every six to eight hours. Exceed 10mg/day with caution. Switch to oral therapy when feasible. IM dose is generally about one-third to one-half the oral dose.</p>	<p>Common: Hypotension, Extrapyramidal disease, Tardive dyskinesia, Nasal Congestion</p> <p>Serious: Agranulocytosis, Leukopenia, Neutropenia, Thrombocytopenia, Drug induced Lupus, Dystonia, Neuroleptic malignant syndrome</p>
<b>Fluphenazine Decanoate (Long Acting Injectable)</b>			
	Schizophrenia, chronic	<p>Initial 12.5-25mg IM or SQ with the dose being repeated or increased as needed and tolerated; generally a single injection may control symptoms up to 4 to 6 weeks.</p> <p>Titration: Dosage should not exceed 100mg. If doses greater than 50mg are deemed necessary, the next dose and succeeding doses should be increased cautiously in increments of 12.5mg</p>	<p>Common: Hypotension, Extrapyramidal Disease, Tardive dyskinesia</p> <p>Serious: QT prolongation, Paralytic ileus, Agranulocytosis, Leukopenia, Thrombocytopenia, Liver Damage, Drug induced Lupus, Neuroleptic Malignant Syndrome</p>
<b>Haloperidol Lactate (Haldol)</b>			
	Schizophrenia	<p>Initial/ Maintenance 0.5-5mg orally 2-3 times a day</p> <p>Injection (acute agitation) 2 to 5 mg IM; may repeat every 4 to 8 hours depending on patient response; increase to every 1 hour if needed; MAX 20mg/day</p>	<p>Common: Hypotension, Constipation, Xerostomia, Extrapyramidal Disease, Somnolence, Blurred Vision</p> <p>Serious: QT prolongation (particularly in IV form), Paralytic ileus, Priapism, Agranulocytosis, Dystonia, Neuroleptic Malignant Syndrome</p>

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b><i>Haloperidol Decanoate (Long Acting Injectable)</i></b>			
	Schizophrenia, chronic	<p>Patients stabilized on oral doses up to 10mg/day 10-15 times previous daily oral dose IM monthly Initial doses &gt;100mg should be administered in 2 separate doses</p> <p>Patients stabilized on oral doses &gt;10mg/day 20 times previous daily oral dose IM for the first month, then 10-15 times previous oral dose IM monthly</p> <p>Max dose Doses &gt;450mg monthly have limited clinical experience</p>	<p>Common: Extrapyramidal disease</p> <p>Serious: QT prolongation, Agranulocytosis, Leukopenia, Anaphylaxis, Neuroleptic Malignant Syndrome, Seizure, Pulmonary Embolism</p>
<b><i>Thiothixene (Navane)</i></b>			
	Schizophrenia	<p>Initial 2 mg orally 3 times daily (milder conditions) or 5mg orally twice daily (severe conditions) with gradual titration to optimal clinical effect.</p> <p>Typical dosage 20-30mg/day</p> <p>Max dose 60mg/day</p>	<p>Common: Orthostatic hypotension, Diminished sweating, Photosensitivity, Constipation, Xerostomia, Extrapyramidal Disease, Somnolence, Blurred vision, Epithelial keratopathy, Retinitis Pigmentosa, Urinary retention</p> <p>Serious: QT prolongation, Paralytic Ileus, Agranulocytosis, Thrombocytopenia, Cholestatic jaundice, Drug induced lupus</p>

# Medications for the Treatment of Bipolar Disorder and Schizophrenia

## SECOND GENERATION ANTIPSYCHOTICS

Aripiprazole (Abilify)			
	Bipolar I disorder-  Acute treatment of manic or mixed episodes; in combination with lithium or valproate; Adjunct  Acute treatment of manic or mixed episodes; monotherapy	Initial 10-15mg orally once a day; target dose 15 mg once a day; may increase to MAX 30 mg/day  Initial and target dose, 15 mg orally once a day; may increase to MAX dose of 30 mg/day	Common: Weight gain, Constipation, Nausea/Vomiting, Extrapyramidal disease, Headache, Insomnia, Sedation, Blurred vision, Anxiety  Serious: QT prolongation, Myocardial Infarction, DKA, Pancreatitis, Agranulocytosis, Leukopenia, CVA/TIA, Suicidal behavior, Angioedema
	Schizophrenia	Initial 10-15mg orally once daily  Titration May increase after 2 weeks at each dose strength	
Aripiprazole Lauroxil (Abilify Maintena)			
	Bipolar I disorder- maintenance monotherapy	Initial/ Maintenance 400mg IM once monthly continue oral aripiprazole after initial injection for 14 consecutive days	Common: Weight gain, Constipation, Nausea/Vomiting, Extrapyramidal disease, Headache, Insomnia, Sedation, Blurred vision, Anxiety  Serious: QT prolongation, Myocardial Infarction, DKA, Pancreatitis, Agranulocytosis, Leukopenia, CVA/TIA, Suicidal behavior, Angioedema
	Schizophrenia	Same as above	
Asenapine (Saphris)			
	Bipolar I disorder – <i>adjunct</i>	Initial 5mg sublingually twice daily with either lithium or valproate  Titration May increase to 10mg twice daily  Max dose Safety of doses >10mg twice daily is not established	Common: Oral Hypoesthesia, Myalgias, Extrapyramidal disease, Dizziness  Serious: QT prolongation, Hyperglycemia, Hypercholesterolemia, Hypertriglyceridemia, Weight gain, Agranulocytosis, Hypersensitivity, Somnolence, Suicidal thoughts
	Bipolar I disorder <i>monotherapy</i>	Initial/ Maintenance 5-10mg sublingually twice daily	
	Schizophrenia	Initial/ Maintenance	



## Medications for the Treatment of Bipolar Disorder and Schizophrenia

		5 mg sublingually twice daily  Titration after 1 week may increase to 10mg twice daily	
<b>Brexipiprazole (Rexulti)</b>			
	Schizophrenia	Initial 1mg orally once daily on days 1-4  Titration 2mg orally once daily on days 5-7  Typical dosage Target dosage of 4mg orally once daily beginning on day 8 based on response and tolerability  Max dose 4mg	Common: Hyperglycemia, Hypertriglyceridemia, Weight gain, Extrapyramidal disease, Headache  Serious: CVA/TIA, Orthostatic hypotension, Syncope, DKA, Agranulocytosis, Seizure, Increased suicidal thoughts, Neuroleptic Malignant Syndrome
<b>Cariprazine (Vraylar)</b>			
	Bipolar I disorder, Acute mixed or manic episodes	Initial 1.5mg orally once daily on day 1 Titration Increase to 3mg once daily on day 2 Further adjust dose in 1.5 or 3mg increments as needed Max dose 6mg/day	Common: Indigestion, Vomiting, Extrapyramidal disease, Somnolence  Serious: Ischemic stroke, Orthostatic hypotension, Diabetes Mellitus, Hyperglycemia, Dyslipidemia, Esophageal dysmotility, Leukopenia, Seizure, Increased Risk of Suicide
	Bipolar I disorder, Maintenance	Maintenance Continue 3 to 6 mg orally once daily	
	Schizophrenia	Initial 1.5mg orally once daily on day 1  Titration Increase to 3mg once daily on day 2 Further adjust dose in 1.5 or 3mg increments as needed  Maintenance 1.5 to 6mg daily  Max dose 6mg/day	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b><i>Clozapine (Clozaril)</i></b>			
	Schizophrenia, Treatment Resistant or with Recurrent Suicidal Behavior	<p>Initial 12.5mg orally 1-2 times daily</p> <p>Titration Increase in increments of 25 to 50mg per day as tolerated</p> <p>Target maintenance dose 300-450mg/day in 2-3 divided doses by the end of 2 weeks</p> <p>Rapid titration in hospitalized patients Day 1-50mg followed by 50-100mg as needed every 6 hours up to a max of 150 additional mg Continue with daily increases in increments of 50-100mg per day thereafter as tolerated.</p> <p>Max dose 900mg/day</p>	<p>Common: Hypotension, Syncope, Tachycardia, Sweating, Weight gain, Constipation, Excess salivation, Nausea, Fever</p> <p>Serious: Cardiomyopathy, QT prolongation, Diabetes Mellitus, Hepatotoxicity, Neuroleptic Malignant Syndrome</p> <p><b>Life Threatening:</b> Agranulocytosis</p> <ul style="list-style-type: none"> <li>– prior to initiation, baseline absolute neutrophil count should be at least 1500/mcL or 1000/mcL for patients with documented ethnic neutropenia</li> <li>– CBC with differential and ANC weekly for the first 6 months, then once every two weeks for the next 6 months, then once a month</li> </ul>
<b><i>Iloperidone (Fanapt)</i></b>			
	Schizophrenia	<p>Initial 1 mg orally twice daily</p> <p>Titration Slow increase not to exceed increase of 2mg twice daily or 4mg/day</p> <p>Target maintenance dose 6-12mg twice daily</p> <p>Max dose 12mg twice daily</p>	<p>Common: Orthostatic hypotension, Tachycardia, Weight gain, Hyperprolactinemia, Xerostomia, Dizziness, Somnolence, Congestion</p> <p>Serious: Hyperglycemia, CVA/TIA, Neuroleptic malignant syndrome</p>
<b><i>Lurasidone (Latuda)</i></b>			
	Bipolar disorder, depressed phase <i>in combination with lithium or valproate</i>	<p>Initial 20mg orally once daily with food (350 cal)</p> <p>Max dose 120mg daily</p>	<p>Common: Dyslipidemia, Hyperglycemia, Weight gain, Diarrhea, Nausea/Vomiting, Extrapyramidal disease, Somnolence</p> <p>Serious: Orthostatic hypotension, Agranulocytosis, CVA/TIA, Suicidal thoughts, Increased creatinine, Neuroleptic Malignant Syndrome</p>
	Schizophrenia	<p>Initial 40mg orally once daily with food (350 cal)</p> <p>Max dose 160mg daily</p>	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b>Quetiapine (Seroquel)</b>			
	Bipolar disorder, depressed phase <i>Monotherapy in acute management</i>	<p>Initial 50mg orally once daily on day 1</p> <p>Titration 100 mg once daily on day 2; 200mg once daily on day 3; 300mg once daily on day 4, all doses given at bedtime</p> <p>Max dose 300mg/day</p>	<p>Common: Orthostatic hypotension, Hypercholesterolemia, Hypertriglyceridemia, Weight Gain, Abdominal pain, Constipation, Xerostomia, Increased LFTs, Weakness, Dizziness, Extrapyramidal disease, Headache, Somnolence</p> <p>Serious: QT prolongation, Syncope, DKA, Pancreatitis, Agranulocytosis, Seizure, Suicidal thoughts, Neuroleptic Malignant Syndrome</p>
	Bipolar disorder, maintenance <i>in combination with lithium or divalproex</i>	<p>Initial/ maintenance 400 to 800mg per day orally divided twice daily, generally continuation of stabilization dose</p> <p>Max dose 800mg/day</p>	
	Bipolar Mania- Adults  <i>Monotherapy <u>or</u> as an adjunct to lithium or divalproex</i>	<p>Initial day 1: 100mg orally divided twice daily</p> <p>Titration day 2: 200mg divided twice daily day 3: 300mg divided twice daily day 4: 400mg divided twice daily; with further dose adjustments in increments of not more than 200mg/day</p> <p>Max dose 800mg/day by day 6</p>	
	Schizophrenia	<p>Initial day 1: 25mg orally twice daily</p> <p>Titration increase total daily dosage by 25 to 50mg divided into 2 to 3 doses on days 2 and 3 to achieve a target dose of 300 to 400mg by day 4 further dose adjustment should generally occur in increments of 20 to 50mg twice daily at intervals of no less than 2 days</p> <p>Max dose 750mg/day</p>	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b>Quetiapine Extended Release (Seroquel)</b>			
	Bipolar disorder, depressed phase <i>Monotherapy in acute management</i>	<p>Initial 50mg orally once daily on day 1</p> <p>Titration 100 mg once daily on day 2; 200mg once daily on day 3; 300mg once daily on day 4, with all doses at bedtime</p> <p>Max dose 300mg/day</p>	<p>Common: Orthostatic hypotension, Hypercholesterolemia, Hypertriglyceridemia, Weight Gain, Abdominal pain, Constipation, Xerostomia, Increased LFTs, Weakness, Dizziness, Extrapyramidal disease, Headache, Somnolence</p> <p>Serious: QT prolongation, Syncope, DKA, Pancreatitis, Agranulocytosis, Seizure, Suicidal thoughts, Neuroleptic Malignant Syndrome</p>
	Bipolar disorder, maintenance <i>in combination with lithium or divalproex</i>	<p>Initial/ maintenance 400 to 800mg per day orally divided twice daily, generally continuation of stabilization dose</p> <p>Max dose 800mg/day</p>	
	Bipolar Mania- Adults  <i>Monotherapy or as an adjunct to lithium or divalproex</i>	<p>Initial day 1: 300mg orally in the evening</p> <p>Titration day 2: 600mg divided twice daily day 3: further dosage adjustment to usual daily dose (400-800mg)</p> <p>Max dose 800mg/day</p>	
	Schizophrenia	<p>Initial 300mg orally in the evening on day 1</p> <p>Titration usual maintenance dose between 400 to 800mg once daily dose increases may occur at intervals of at least 1 day in increments of up to 300mg/day</p> <p>Max dose 800mg/day</p>	
	Schizophrenia, Maintenance	400 to 800mg per day orally once daily in the evening	
<b>Paliperidone (Invega)</b>			
	Bipolar I disorder, Acute manic and mixed episodes <i>(off label)</i>	<p>Initial 6mg orally once daily</p> <p>Titration</p>	<p>Common: Tachycardia, Hyperprolactinemia, Weight gain, Constipation, Indigestion, Extrapyramidal Disease, Somnolence, Tremor</p>

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

		Adjust based on efficacy and tolerance with 3mg/day increments at intervals of more than 4 days  Max dose 12mg/day	Serious: QT prolongation, Agranulocytosis, Dysphagia, Tardive dyskinesia
	Schizoaffective disorder	Same as above	
	Schizophrenia	Initial 6mg orally once daily  Titration Adjust based on efficacy and tolerance with 3mg/day increments at intervals of more than 5 days  Max dose 12mg/day	
Ziprasidone (Geodon)			
	Bipolar I disorder, Acute manic or mixed episodes, <i>monotherapy</i>	Initial 40mg twice daily with food on day 1  Titration 60 or 80mg twice daily with food on day 2 Adjust based on tolerance and efficacy within range of 40 to 80mg twice daily  Max dose Up to 80mg twice a day	Common: Rash, Weight gain, Constipation, Nausea/Vomiting, Headache, Somnolence, Extrapyramidal disease, Dizziness  Serious: Dysphagia, QT prolongation, Seizure, Neuroleptic malignant syndrome
	Bipolar I disorder, <i>adjunct with lithium or valproate</i>	Initial/ maintenance 40 to 80mg orally twice daily with food	
	Schizophrenia	Initial 20mg orally twice a day with food  Titration May gradually increase dosage not less than every 2 days  Max dose Up to 80mg twice a day  Periodically reassess the need for maintenance treatment and use lowest effective dose; no additional benefit has been demonstrated for doses greater than 20mg twice daily	

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b>Risperidone (Risperdal)</b>			
	Bipolar I disorder	<p>Initial 2-3mg orally once a day</p> <p>Maintenance Dosage adjustments should be made in increments of 1mg/day at intervals of at least 24 hours,</p> <p>Max dose Doses higher than 6mg/day have not been evaluated in clinical trials</p>	<p>Common: Rash, Hyperprolactinemia, Weight gain, Constipation, Diarrhea, Excess salivation, Nausea/Vomiting, Xerostomia, Extrapyramidal Disease, Sedation, Tremor, Anxiety</p> <p>Serious: QT prolongation, Syncope, DKA, Pancreatitis, Agranulocytosis, Thrombocytopenia, Seizure, Priapism</p>
	Schizophrenia	<p>Initial 2mg/day orally, administered in either one or two doses</p> <p>Titration Increase as tolerated in increments of 1 to 2 mg/day at intervals not less than 24 hours</p> <p>Typical dosage 4 to 8mg/day</p> <p>Max dose Doses higher than 16mg/day have not been evaluated in clinical trials</p>	<p>Common: Rash, Hyperprolactinemia, Weight gain, Constipation, Diarrhea, Excess salivation, Nausea/Vomiting, Xerostomia, Extrapyramidal Disease, Sedation, Tremor, Anxiety</p> <p>Serious: QT prolongation, Syncope, DKA, Pancreatitis, Agranulocytosis, Thrombocytopenia, Seizure, Priapism</p>

## Medications for the Treatment of Bipolar Disorder and Schizophrenia

<b>Risperidone Long Acting Injectable</b> <i>Establish tolerability to oral risperidone prior to initiation of treatment for all indications</i>			
	Bipolar I disorder	<p><b>Initial</b> 25mg IM every 2 weeks, oral risperidone or another antipsychotic medication should be given with the initial injection, continued for 3 weeks, and then discontinued</p> <p><b>Titration</b> Dose may be increased to 37.5mg or 50mg IM with adjustments made not more frequently than every 4 weeks; clinical effects of dose adjustment should not be expected earlier than 3 weeks after any dose change</p> <p><b>Max dose</b> 50mg IM every 2 weeks</p>	<p><b>Common:</b> Rash, Hyperprolactinemia, Weight gain, Constipation, Diarrhea, Excess salivation, Nausea/Vomiting, Xerostomia, Extrapyramidal Disease, Sedation, Tremor, Anxiety</p> <p><b>Serious:</b>  QT prolongation, Syncope, DKA, Pancreatitis, Agranulocytosis, Thrombocytopenia, Seizure, Priapism</p>
	Schizophrenia	<p><b>IM:</b> same as above</p> <p><b>SQ:</b> 90 or 120mg SQ in abdomen once a month, max 90 to 120mg/month; no loading dose or supplemental oral risperidone is advised</p>	

### References:

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T908658, Valproic Acid; [updated 2018 Nov 06]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T908658>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233472, Lamotrigine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233472>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233111, Carbamazepine; [updated 2018 Nov 30]. Available from <https://www.dynamed.com/topics/dmp~AN~T233111>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233030, Lithium carbonate; [updated 2018 Nov 06]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233030>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T356386, Chlorpromazine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T356386>. Registration and login required.

# Medications for the Treatment of Bipolar Disorder and Schizophrenia

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T908070, Thioridazine; [updated 2018 Nov 06]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T908070>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T356425, Loxapine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T356425>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233091, Perphenazine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233091>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T907681, Fluphenazine Decanoate; [updated 2018 Nov 06]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T907681>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T907728, Haloperidol; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T907728>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T356499, Thiothixene; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T356499>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T232892, Aripiprazole; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T232892>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T900407, Asenapine; [updated 2018 Nov 30] Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T900407>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T909096, Brexpiprazole; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T909096>. Registration and login required

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233402, Clozapine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233402>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T900405, Iloperidone; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T900405>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T900696, Lurasidone; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T900696>. Registration and login required.



# Medications for the Treatment of Bipolar Disorder and Schizophrenia

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233518, Olanzapine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233518>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233557, Quetiapine; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233557>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T356351, Paliperidone; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T356351>. Registration and login required

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T908295, Ziprasidone Hydrochloride; [updated 2018 Nov 06]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T908295>. Registration and login required.

DynaMed [Internet]. Ipswich (MA): EBSCO Information Services. 1995 - . Record No. T233460, Risperidone; [updated 2018 Nov 30]. Available from <https://www.dynamed-com.evms.idm.oclc.org/topics/dmp~AN~T233460>. Registration and login required.

Jibson MD, Marder S, First-generation antipsychotic medications: Pharmacology, administration, and comparative side effects Solomon D, ed. In: *UpToDate*, Waltham Mass.: UpToDate, 2018.

Jibson MD, Marder S, Second-generation antipsychotic medications: Pharmacology, administration, and side effects Solomon D, ed. In: *UpToDate*, Waltham Mass.: UpToDate, 2020.

## Resources

DailyMed: <https://dailymed.nlm.nih.gov>

Micromedex: <https://www.micromedexsolutions.com>

Lexicomp: <https://online.lexi.com>

## **Resources**

National Institute of Mental Health (NIMH) <https://nimh.nih.gov/health/index.shtml>

National Alliance on Mental Illness (NAMI) <https://nami.org/About-Mental-Illness>

Substance Abuse and Mental Health Services Administration (SAMHSA) <https://www.samhsa.gov>

Depression and Bipolar Support Alliance <https://dbsalliance.org>

Mental Health America <https://mhanational.org>

SMI Adviser <https://www.smiadviser.org>

## **Additional Reference**

**The American Psychiatric Association Practice Guideline for the Treatment of Patients with Schizophrenia, Third Edition.**  
**Published September 2020, Copyright 2021.** [psychiatryonline.org/doi/epdf/10.1176/appi.books.9780890424841](https://psychiatryonline.org/doi/epdf/10.1176/appi.books.9780890424841)