# SENTARA HEALTH PLANS

### PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

# Migraine Treatment: Injectable Calcitonin Gene-Related Peptide (CGRP) **Antagonists**

ow)
PREFERRED
□ Emgality® (galcanezumab)
NON-PREFERRED
must have tried and failed <u>BOTH</u> preferred agents and meet all
ORMATION: Authorization may be delayed if incomplete.
Date of Birth:
Date:
Fax Number:
ation may be delayed if incomplete.
Length of Therapy:
ICD Code, if applicable:
Date weight obtained:
eviously prescribed injectable calcitonin gene-related peptide (CGRP) equested medication?
□ Yes <b>OR</b> □ No

(Continued on next page)

(Continued from previous page)

•	• If yes, please list the medication that will be discontinued and the medication that will be initiated upon approval along with the corresponding effective date.			
	Medication to be discontinued:	Effective date:	_	
	Madication to be initiated:	Effective date:		

#### **Recommended Dosing & Quantity Limits:**

Drug	Dose	Quantity Limit
Aimovig® (erenumab)	• Migraine Prophylaxis: Initial: 70 mg SC once a month; some members may benefit from 140 mg once a month (given as 2 consecutive 70 mg injections)	<ul> <li>70 mg/mL (1 mL/30 day)</li> <li>140 mg dose (2 mL/30 days)</li> <li>If using the 140 mg dose, must use the package labeled specifically for 140 mg/mL</li> </ul>
Ajovy® (fremanezumab)	<ul> <li>Migraine Prophylaxis for adults 18 years of age or older: 225 mg SC monthly or 675 mg every 3 months</li> <li>Migraine Prophylaxis for children ≥6 years and Adolescents ≤17 years, weighing ≥45 kg: 225 mg SC monthly</li> </ul>	<ul> <li>225 mg/1.5 mL; 1.5 mL (1</li> <li>syringe) per 30 days or 4.5 mL (3 syringes) per 90 days</li> </ul>
Emgality® (galcanezumab)	<ul> <li>Migraine Prophylaxis: Initial: 240 mg SC as a single loading dose, followed by 120 mg once monthly</li> <li>Episodic cluster headache prophylaxis: 300 mg SC at the onset of the cluster period and then once monthly until the end of the cluster period</li> </ul>	<ul> <li>120 mg/mL; 1 mL (1 auto-injector and prefilled syringe) per 30 days with one time loading dose of 2 mL (2 auto-injectors)</li> <li>For Episodic Cluster headache diagnosis only: 300 mg dose; 100 mg/mL prefilled syringe</li> </ul>

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

### **Authorization Criteria**

	Member must	meet <u>ONE</u> o	of the	following	age requirement	s:
--	-------------	-------------------	--------	-----------	-----------------	----

- □ For Aimovig & Emgality requests: Member must be 18 years of age or older
- ☐ For Ajovy requests: Member must be 6 years of age or older and weigh 45 kg or more
- □ Provider has attested to all clinical criteria for <u>ONE</u> of the applicable diagnoses below

(Continued on next page)

DI	[A	GN	OSI	S:	Please check <b>ONE</b> of the applicable diagnoses below
		hro qua		&	Episodic Migraine Headache Prevention (All applicable boxes below must be met
		follo	owir Mer Mer prop treat evid	ng: nbe nbe bhy me enc An Bet An An	ticonvulsants (divalproex, valproate, topiramate) a blockers (atenolol, metoprolol, nadolol, propranolol, timolol) tidepressants (amitriptyline, venlafaxine) giotensin II receptor blocker (candesartan) *requires prior authorization* ectable CGRP inhibitors (Aimovig®, Emgality®, Ajovy®) or oral CGRP inhibitors indicated
		pref	·Ajo	vy	migraine prevention (Qulipta <sup>TM</sup> , Nurtec ODT <sup>®</sup> ) *requires prior authorization*  ® requests for members $\geq 18$ years of age: Member must have tried and failed <b>BOTH</b> gents Aimovig® and Emgality® <b>AND</b> meet all prior authorization criteria for approval of
		Rec	ques abot	ulir	For concurrent use of Calcitonin Gene-Related Peptide (CGRP) inhibitors with Botox® numtoxinA) for migraine headache prevention (if applicable): Member must meet <u>ALL</u> the criteria (verified by chart notes and/or pharmacy paid claims):
			Mer	nbe	er must have a diagnosis of Chronic or Episodic Migraine Headache and is continuing to ence $\geq$ 4 migraine headache days per month after receiving therapy with <u>ALL</u> the following
			1 • 1	mig of I nig	mber must have failed a <b>2-month</b> trial of at least one medication from <b>TWO</b> different graine prophylactic classes supported by the American Headache Society/American Academy Neurology treatment guidelines 2012/2015/2021/2024, Level A and B evidence: ICSI 2013, h quality evidence:  Anticonvulsants (divalproex, valproate, topiramate)  Beta blockers (atenolol, metoprolol, nadolol, propranolol, timolol)
			[	_	Antidepressants (amitriptyline, venlafaxine)
				<b>_</b>	Angiotensin II receptor blocker (candesartan) *requires prior authorization*
					mber must meet <u>ONE</u> of the following:
			[		Member has had an inadequate response to a <u>2-month</u> trial with an injectable CGRP inhibitor (e.g., Aimovig <sup>®</sup> , Ajovy <sup>®</sup> , Emgality <sup>®</sup> ) or an oral CGRP inhibitor indicated for migraine prevention (e.g., Nurtec <sup>®</sup> ODT, Qulipta <sup>™</sup> ) *requires prior authorization*
			[		Member has had an inadequate response to a <u>6-month</u> trial (2 injection cycles) of Botox <sup>®</sup> (onabotulinumtoxinA) *requires prior authorization*

(Continued on next page)

#### PA Migraine Treatment: Injectable CGRP Antagonists (CORE) (Continued from previous page)

Episodic Cluster Headaches (Emgality® Only) (All applicable boxes below must be met to qualify)
Member has between one headache every other day and eight headaches per day
Member must have failed at least a <u>1-month</u> trial of at least <u>ONE</u> generic standard prophylactic pharmacologic therapy, used to prevent cluster headache and supported by the American Headache Society/American Academy of Neurology treatment guidelines:
□ Suboccipital steroid injection

□ Calcium channel blockers (verapamil) ☐ Alkali metal/ Antimanic (lithium)

☐ Anticoagulant (warfarin)

☐ Anticonvulsants (topiramate)

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. \*