SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: **The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request**. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Benlysta® (belimumab) **Subcutaneous Injection (Pharmacy)**

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ **Date of Birth:** _____

Prescriber Name: _____

Prescriber Signature: _____ **Date:** _____

Office Contact Name: _____

Phone Number: _____ **Fax Number:** _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ **Length of Therapy**: _____

Diagnosis: _____ **ICD Code, if applicable:** _____

Weight (if applicable): _____ **Date weight obtained:** _____

Recommended Dosing:

Diagnosis	Adults (Auto-injector or Prefilled syringe)	Pediatric Patients 5 to less than 18 years of age (Auto-injector only)
SLE	200 mg once weekly	• Patients ≥ 40 kg: 200 mg once weekly • Patients 15 kg to <40 kg: 200 mg once every 2 weeks
Lupus Nephritis	400 mg once weekly x 4 doses, followed by 200 mg once weekly	• Patients ≥ 40 kg: 400 mg once weekly for 4 doses, followed by 200 mg once weekly • Patients 15 kg to <40 kg: 200 mg once weekly for 4 doses, followed by 200 mg once every 2 weeks

Quantity Limits: 200 mg once weekly (4 injections per 28 days)

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

❑ **Diagnosis - active systemic lupus erythematosus (SLE) in patients who are receiving standard therapy**

Initial Authorization: 12 months

- ❑ Prescribed by or in consultation with a rheumatologist
- ❑ Member is 5 years of age or older with a diagnosis of active, autoantibody-positive SLE confirmed by **ONE** of the following **(submit lab results)**:
  - ❑ anti-nuclear antibody (ANA) titer ≥ 1:80
  - ❑ anti-double stranded DNA (anti-dsDNA) ≥ 30 IU/mL
- ❑ Member's SLE activity has been confirmed by **ONE** of the following **(submit results)**:
  - ❑ Safety of Estrogen in Lupus National Assessment – Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI) score of 6-12
  - ❑ ≥2 British Isles Lupus Assessment Group (BILAG) B organ domain scores
- ❑ Member has tried three of the following and is established on two of the following therapies taken for the last 90 days **(please submit chart notes documenting therapy trials with insufficient disease control)**:

❑ mycophenolate	❑ hydroxychloroquine	❑ azathioprine
❑ cyclophosphamide	❑ methotrexate	❑ cyclosporine
❑ corticosteroids	❑ Other_____	

- ❑ Member does **NOT** have any of the following limitations to therapy: severe active central nervous system lupus, current or previous diagnosis of progressive multifocal leukoencephalopathy (PML), or concurrent use with other biologics

❑ **Diagnosis - active systemic lupus erythematosus (SLE) in patients who are receiving standard therapy**

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ❑ All initial authorization criteria continues to be met

(Continued on next page)

- ❑ Member's response to therapy has been confirmed by **ONE** of the following **(submit results)**:
  - ❑ Safety of Estrogen in Lupus National Assessment – Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI) score has improved by and/or maintained at a level that is ≥4 points below baseline score
  - ❑ No new BILAG-A organ domain score OR 2 new BILAG-B organ domain scores
- ❑ Member has an absence of intolerable side effects such as serious infections, signs or symptoms of progressive multifocal leukoencephalopathy (PML), malignancy, severe hypersensitivity reactions/anaphylaxis, or serious infusion reactions

❑ Diagnosis - active lupus nephritis in patients who are receiving standard therapy

Initial Authorization: 12 months

- ❑ Prescribed by or in consultation with a nephrologist or rheumatologist
- ❑ Member is 5 years of age or older with a diagnosis of active lupus nephritis Class III, IV, or V as confirmed by renal biopsy
- ❑ Member's diagnosis of active, autoantibody-positive SLE was confirmed by **ONE** of the following **(submit lab results)**:
  - ❑ anti-nuclear antibody (ANA) titer ≥ 1:80
  - ❑ anti-double stranded DNA (anti-dsDNA) ≥ 30 IU/mL
- ❑ Member has active renal disease and has received standard therapy for the last 90 days with corticosteroids along with **ONE** of the following **(chart notes documenting established therapy must be submitted)**:
  - ❑ mycophenolate
  - ❑ cyclophosphamide
- ❑ Provider must obtain a baseline measurement of **ONE** of the following collected within the last 30 days **(labs must be submitted)**:
  - ❑ urine protein:creatinine ratio (uPCR)
  - ❑ urine protein
- ❑ Member does **NOT** have any of the following limitations to therapy: severe active central nervous system lupus, current or previous diagnosis of progressive multifocal leukoencephalopathy (PML), or concurrent use with other biologics

❑ Diagnosis - active lupus nephritis in patients who are receiving standard therapy

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ❑ All initial authorization criteria continues to be met

(Continued on next page)

- ❑ Member has had improvement from baseline and/or stabilization since last approval of one of the following **(submit current labs completed within the last 30 days)**:
  - ❑ urine protein:creatinine ratio (uPCR)
  - ❑ urine protein
- ❑ Member has an absence of intolerable side effects such as serious infections, signs or symptoms of progressive multifocal leukoencephalopathy (PML), malignancy, severe hypersensitivity reactions/anaphylaxis, or serious infusion reactions

Medication being provided by a Specialty Pharmacy – Proprium Rx

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.