

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

**Drug Requested:** Kerendia<sup>®</sup> (finerenone)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

**Quantity Limit:** 30 tablets per 30 days

### **Kerendia<sup>®</sup> Initial Dosing Recommendations:**

eGFR (mL/min/1.73m <sup>2</sup> )	Starting Dose
≥ 60	20 mg once daily
≥ 25 to < 60	10 mg once daily
< 25	Not Recommended

(Continued on next page)

<b>Dose Adjustment Based on Current Serum Potassium Concentration and Current Kerendia® Dose</b>			
		<b>10 mg once daily</b>	<b>20 mg once daily</b>
<b>Current Serum Potassium (mEq/L)</b>	<b>≤ 4.8</b>	Increase the dose to 20 mg once daily*	Maintain 20 mg once daily
	<b>&gt; 4.8 – 5.5</b>	Maintain 10 mg once daily	Maintain 20 mg once daily
	<b>&gt; 5.5</b>	Withhold Kerendia®. Consider restarting at 10 mg once daily when serum potassium ≤ 5.0 mEq/L	Withhold Kerendia®. Restart at 10 mg once daily when serum potassium ≤ 5.0 mEq/L

\*If eGFR has decreased by more than 30% compared to previous measurement, maintain 10 mg dose.

**CLINICAL CRITERIA:** Check below all that apply. **All criteria must be met for approval.** To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial Authorization: 6 months**

- Member is 18 years of age or older and has a diagnosis of chronic kidney disease associated with type 2 diabetes
- Must submit lab test results documenting **BOTH** of the following obtained within the past 60 days
  - Members' current eGFR is > 25mL/minute/1.73 m<sup>2</sup>
  - Member's current Urinary Albumin-to -Creatinine Ratio (UACR) is ≥ 30 mg/g
- Member's current serum potassium is ≤5 mEq/L along with **BOTH** of the following (**submit current lab documentation obtained within the past 60 days**):
  - Therapy will **NOT** be initiated if serum potassium >5 mEq/L
  - Initiation with increased serum potassium monitoring during the first 4 weeks will be preformed if serum potassium is >4.8 to 5 mEq/L
- Member is established on treatment with maximally tolerated dose of angiotensin-converting enzyme inhibitor (ACE) or angiotensin receptor blocker (ARB) medication and will continue to take along with Kerendia® (finerenone)
- Member tried and failed, has a contradiction, or intolerance to at least **ONE** sodium glucose transport protein 2 (SGLT2) inhibitor that is indicated for use in patients with chronic kidney disease (e.g., Farxiga®, Jardiance®)
- Member doses **NOT** have a diagnosis of adrenal insufficiency or a diagnosis of know significant non-diabetic renal disease, including clinically relevant renal artery stenosis
- Member is **NOT** receiving simultaneous treatment with strong CYP3A4 inhibitors
- For initial therapy, member will be dosed as follows:
  - eGFR ≥60 mL/minute/1.73 m<sup>2</sup>: starting dose will be 20 mg once daily
  - eGFR ≥25 to <60 mL/minute/1.73 m<sup>2</sup>: starting dose will be 10 mg once daily

(Continued on next page)

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member's current eGFR is  $>25$  mL/minute/1.73 m<sup>2</sup> (**submit current lab documentation**)
- Member's Urinary Albumin-to-creatinine Ratio (UACR) has decreased by  $\geq 30\%$  from baseline level and/or been sustained at  $\geq 30\%$  below baseline level since last approval (**submit current lab documentation**)
- Member's current serum potassium level does **NOT** exceed 5.5 mEq/L (**submit current lab documentation**)
- Provider attests Kerendia<sup>®</sup> will be withheld if serum potassium is  $>5.5$  mEq/L and will consider restarting therapy when serum potassium normalizes ( $\leq 5.0$  mEq/L)

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****