

# SENTARA HEALTH PLANS

## MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-668-1550**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

**For Medicare Members:** Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Additional indications may be covered at the discretion of the health plan.

**Drug Requested: Kebilidi (eladocogene exuparvec-tneq) (J3590) (Medical)**

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

### **Dosing Limits**

#### **A. Quantity Limit (max daily dose) [NDC Unit]:**

- Kebilidi is supplied in a single-dose vial that contains  $2.8 \times 10^{11}$  vg of eladocogene exuparvec-tneq in an extractable volume of 0.5 mL of suspension. Each mL of suspension contains  $5.6 \times 10^{11}$  vg of eladocogene exuparvec-tneq [NDC 52856-0601-XX]

(Continued on next page)

**B. Max Units (per dose and over time) [HCPCS Unit]:**

- One treatment (dose) per lifetime.
- Administer a total dose of  $1.8 \times 10^{11}$  vg (0.32 mL total volume) delivered as four 0.08 mL ( $0.45 \times 10^{11}$  vg) infusions (two sites per putamen-anterior and posterior) at a rate of 0.003 mL/minute (0.18 mL/hour) for a total of 27 minutes per site, administered in a single stereotactic surgery using a cannula that is FDA-authorized for intraparenchymal infusion.

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Length of Authorization: 3 months with an allowance of only 1 dose per lifetime**

**Coverage will be provided for one treatment course and may NOT be renewed.**

- Member is at least 16 months of age through 10 years of age
- Prescribed by or in consultation with a pediatric neurologist, neurologist or neurosurgeon
- Member has biallelic pathogenic variants in the dopa decarboxylase (DDC) gene (**submit documentation**)
- Member has decreased aromatic L-amino acid decarboxylase (AADC) enzyme activity in plasma per current laboratory standards (**submit documentation**)
- According to the prescribing physician, the patient has continued symptoms of AADC deficiency despite use of at least one standard medication therapy [**Note: Examples of medications used for AADC deficiency include dopamine agonists (e.g., pramipexole, ropinirole, rotigotine), monoamine oxidase inhibitors (e.g., tranlycypromine, selegiline), pyridoxine, and other forms of vitamin B6**]
- Member is unable to ambulate independently
- Member has achieved skull maturity as assessed by neuroimaging
- Member does **NOT** have pyridoxine 5'-phosphate oxidase or tetrahydrobiopterin (BH4) deficiency
- Member has **NOT** received prior gene therapy or Kebilidi in the past (**verified by medical paid claims**) [**NOTE:** Verify through claims history that the member has not previously received any other gene therapy or Kebilidi. If no claim for Kebilidi is present (or if claims history is not available), the prescribing physician confirms that the member has not previously received any other gene therapy or Kebilidi.]
- Member must **NOT** have a baseline anti-AAV2 antibody titer above 1:1200 or >1 optical density value by enzyme-linked immunosorbent assay
- Member does **NOT** have any contraindications that would preclude surgical intra putaminal administration
- Member has tested negative for coronavirus disease of 2019 (COVID-19) a maximum of 72 hours prior to receiving gene therapy

(Continued on next page)

**Medication being provided by: Please check applicable box below.**

- Location/site of drug administration: \_\_\_\_\_  
NPI or DEA # of administering location: \_\_\_\_\_

**OR**

- Specialty Pharmacy

For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****  
***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****