

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Trikafta® (elexacaftor/tezacaftor/ivacaftor and ivacaftor)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Recommended Dosing:

- **Children ≥ 2 years to < 6 years weighing < 14 kg:** Oral: 1 packet (containing elexacaftor 80 mg/tezacaftor 40 mg/ivacaftor 60 mg) in the morning and 1 packet (containing ivacaftor 59.5 mg) in the evening
- **Children ≥ 2 years to < 6 years weighing > 14 kg:** Oral: 1 packet (containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and 1 packet (containing ivacaftor 75 mg) in the evening
- **Children ≥ 6 years to < 12 years weighing < 30 kg:** Oral: 2 tablets (each containing elexacaftor 50 mg/tezacaftor 25 mg/ivacaftor 37.5 mg) in the morning and ivacaftor 75 mg in the evening, approximately 12 hours apart
- **Children ≥ 6 years to < 12 years weighing ≥ 30 kg:** Oral: 2 tablets (each containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and ivacaftor 150 mg in the evening, approximately 12 hours apart
- **Children ≥ 12 years, Adolescents and Adults:** Oral: 2 tablets (each containing elexacaftor 100 mg/tezacaftor 50 mg/ivacaftor 75 mg) in the morning and one ivacaftor 150 mg tablet in the evening, approximately 12 hours apart

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- ☐ Member is **2 years of age or older** with a diagnosis of Cystic Fibrosis
- ☐ Member has **at least one** of the F508del mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene as confirmed by an FDA-cleared test that is responsive to elexacaftor/tezacaftor/ivacaftor (**test results must be attached**)
- ☐ Prescribing physician is a pulmonologist or has consulted with a pulmonologist who specializes in the treatment of Cystic Fibrosis
- ☐ Baseline FEV₁ within the last 30 days must be submitted (**test results must be attached**), unless the member is unable to perform a pulmonary function test (**documentation required**)
- ☐ Number of pulmonary exacerbations or hospitalizations in the preceding 6 months must be noted:

- ☐ Baseline body mass index must be noted: _____
- ☐ Baseline liver function tests have been completed prior to initiating therapy and will be completed annually (**labs must be attached**)
- ☐ Provider attests a baseline ophthalmic examination to monitor lens opacities/cataracts has been completed for pediatric members
- ☐ Member does **NOT** have severe hepatic impairment (Child-Pugh Class C)
- ☐ Member will **NOT** be taking Trikafta® in combination with any other CFTR modulator therapy (i.e., Symdeko®, Orkambi®, Kalydeco®, Alyftrek™) **NOTE:** Concurrent therapy with these agents will **NOT** be approved
- ☐ Member will avoid concomitant use of strong CYP3A inducers (e.g., rifampin, carbamazepine, phenytoin, phenobarbital, St. John's wort) and strong or moderate CYP3A inhibitors (e.g., fluconazole, itraconazole)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member continues to meet all initial authorization criteria
- ☐ Member has demonstrated disease response as indicated by **one or more** of the following (**must submit current labs and chart notes**):
 - ☐ Decreased pulmonary exacerbations or hospitalizations compared to pretreatment baseline
 - ☐ Stabilization of lung function as measured by FEV₁ within the last year compared to baseline
 - ☐ Improvement in quality of life, weight gain, or growth

(Continued on next page)

- ☐ Member has **NOT** received a lung transplant
- ☐ Member has experienced an absence of unacceptable toxicity from therapy (i.e., elevated transaminases (ALT or AST), development of cataracts or lens opacities)

Date of initiation of Trikafta [®] therapy: _____	Reauthorization Date: _____
Baseline FEV1 (last FEV1 prior to starting Trikafta [®]): _____	Current FEV1 (FEV1 <u>AFTER</u> last dose of Trikafta [®]): _____
Baseline Weight: _____	Current weight: _____
Baseline BMI: _____	Current BMI: _____
Number of hospitalizations since last approval of Trikafta [®] must be noted _____	

Medication being provided by a Specialty Pharmacy – Proprium Rx
--

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****