

# SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** Panretin® Gel (alitretinoin)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

**Recommended Dosing:** Initial: Apply gel twice daily to cutaneous lesions; may gradually increase application frequency to 3 or 4 times daily based on lesion tolerance. Continue as long as deriving clinical benefit; response may be observed within 2 weeks of initiation; however, most patients require a longer period, and further benefit may be attained with a longer application period (>14 weeks) in some patients.

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial Authorization: 12 months**

- ☐ Member is 18 years of age or older
- ☐ Medication is prescribed by or in consultation with a dermatologist, oncologist, or infectious disease specialist
- ☐ Member has been diagnosed with cutaneous lesions in AIDS-related Kaposi's sarcoma (KS)

(Continued on next page)

- ☐ Member must **NOT** have any of the following exclusions from therapy:
  - Receiving systemic therapy for Kaposi sarcoma
  - Diagnosed with advanced cutaneous, oral visceral, or nodal disease
  - More than 10 new KS lesions in the previous month, symptomatic lymphedema, or symptomatic pulmonary KS

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member continues to derive a positive clinical response treatment
- ☐ Member must **NOT** have any of the following exclusions from therapy:
  - Receiving systemic therapy for Kaposi sarcoma
  - Diagnosed with advanced cutaneous, oral visceral, or nodal disease
  - More than 10 new KS lesions in the previous month, symptomatic lymphedema, or symptomatic pulmonary KS
- ☐ Member has experienced an absence of unacceptable toxicity from the drug (e.g., grade 3 dermal irritation)

**Medication being provided by Specialty Pharmacy – Proprium Rx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****  
***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****